

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 August 2005 (04.08.2005)

PCT

(10) International Publication Number
WO 2005/071068 A1

(51) International Patent Classification⁷: **C12N 7/04**,
A61K 39/12, C12N 15/62, C07K 14/08, 19/00, C12N
15/87, 15/86, 5/10

117, E-28006 Madrid (ES). **RODRÍGUEZ FERNÁN-
DEZ-ALBA, Juan Ramón** [ES/ES]; Ronda de Poniente,
4 - 2° C-D, E-28760 Tres Cantos - Madrid (ES).

(21) International Application Number:
PCT/EP2005/000694

(74) Agent: **ARIAS SANZ, Juan**; ABG Patentes, S.L., Orense,
68, 7th floor, E-28020 Madrid (ES).

(22) International Filing Date: 21 January 2005 (21.01.2005)

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
P200400121 21 January 2004 (21.01.2004) ES

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (for all designated States except US):
**CONSEJO SUPERIOR DE INVESTIGACIONES
CIENTÍFICAS** [ES/ES]; Serrano, 117, E-28006 Madrid
(ES). **BIONOSTRA, S.L.** [ES/ES]; Ronda de Poniente, 4
- 2° C-D, E-28760 Tres Cantos - Madrid (ES).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **RODRÍGUEZ
AGUIRRE, José Francisco** [ES/ES]; Serrano, 117,
E-28006 Madrid (ES). **RUIZ CASTÓN, José** [ES/ES];
Serrano, 117, E-28006 Madrid (ES). **GONZÁLEZ DE
LLANO, María Dolores** [ES/ES]; Serrano, 117, E-28006
Madrid (ES). **OÑA BLANCO, Ana María** [ES/ES];
Serrano, 117, E-28006 Madrid (ES). **ABAITUA ELUS-
TONDO, Fernando** [ES/ES]; Serrano, 117, E-28006
Madrid (ES). **LUQUE BUZO, Daniel** [ES/ES]; Serrano,

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: EMPTY CAPSIDS (VLPs(-VP4)) OF THE INFECTIOUS BURSAL DISEASE VIRUS (IBDV), OBTAINMENT
PROCESS AND APPLICATIONS

(57) Abstract: The empty capsids of the infectious bursal disease virus (IBDV), VLP(-VP4), are characterized in that they are
constituted only by assembly of IBDV pVP2 proteins and IBDV VP3 proteins. Said capsids have immunogenic activity and can be
used in the manufacture of vaccines for protecting animals from the infection caused by IBDV, as well as in the manufacture of gene
therapy vectors.

WO 2005/071068 A1

Best Available Copy

EMPTY CAPSIDS (VLPs(-VP4)) OF THE INFECTIOUS BURSAL DISEASE VIRUS (IBDV), OBTAINMENT PROCESS AND APPLICATIONS

FIELD OF THE INVENTION

5 The invention is related to empty viral capsids of the infectious bursal disease virus (IBDV), with immunogenic activity against IBDV, constituted by the IBDV VP3 and pVP2 proteins, to the production thereof by means of genetic engineering, and to their applications, particularly in the manufacture of vaccines against the avian disease called infectious bursal disease caused by IBDV, and in the manufacture of gene therapy vectors.

10

BACKGROUND OF THE INVENTION

 The infectious bursal disease virus (IBDV), also known as Gumboro disease, belongs to the *Birnaviridae* family, infects different bird species and is directly responsible for a severe immunosuppressive disease causing important economic losses in the world poultry
15 industry.

 IBDV particles are icosahedral, with T=13 symmetry, they lack an envelope and are formed by a single protein layer. Up until now, the approaches aimed at obtaining an atomic model for IBDV particles have failed. As a result, the structural information available is based on three-dimensional models generated from images obtained by electron
20 cryomicroscopy of the purified virus and of the VLPs. Based on these studies, it has been verified that the outer surface of the particle is formed by a continuous lattice of 260 trimers of the VP2 protein (37 kDa) organized in five different formations. The inner face of the particles contains 200 trimers of the VP3 protein (29 kDa), the latter, independent from one another, are bound to the basal area of the VP2 trimers. It has been suggested that a third
25 polypeptide, VP4 (28 kDa), could also be part of the particles, being located at the base of the pentamers forming the vertices of the icosahedral structure.

 The VP2, VP3 and VP4 polypeptides are produced from the proteolytic processing of a polypeptide precursor of a size of 109 kDa. This precursor is auto-catalytically processed, releasing the pVP2 (48 kDa), VP3 and VP4 polypeptides. The VP4 domain, which is located
30 in the central region of the polyprotein, belongs to the Lon protease family and is responsible for the proteolytic cleavage. The pVP2 and VP3 polypeptides are directly responsible for the capsid assembly. The pVP2 product undergoes a last cleavage at its C-terminal end before giving rise to the mature form of the protein, VP2, which is the one found in purified

particles (Da Costa, B., Chevalier, C., Henry, C., Huet, J. C., Petit, S., Lepault, J., Boot, H. & Delmas, B. (2002). The capsid of infectious bursal disease virus contains several small peptides arising from the maturation process of pVP2. *Journal of Virology* 76:2393-2402). This pVP2 processing is necessary for the correct formation of the capsids and requires the presence of VP3, although the responsible protease has not yet been identified (Maraver, A., Oña, A., Abaitua, F., González, D., Clemente, R., Diaz-Ruiz, A., Caston, J. R., Pazos, F. & Rodríguez, J. F. (2003). The oligomerization domain of VP3, the scaffolding protein of infectious bursal disease virus, plays a critical role for capsid formation. *Journal of Virology* 77:6438-49).

Conventional vaccines used for the control of infectious bursal disease are based on the use of strains, with different degrees of virulence, of IBDV itself grown in cell culture or in embryonated eggs. The extracts containing the infectious material are subjected to chemical inactivation processes to produce inactivated vaccines, or else are used directly to produce live attenuated vaccines (Sharma, J. M., Kim, I. J., Rautenschlein, S. & Yeh, H. Y. (2000). Infectious bursal disease virus of chickens: pathogenesis and immunosuppression. *Developmental and Comparative Immunology* 24:223-235; van den Berg TP, Eterradossi N, Toquin D, Meulemans G. 2000. *Rev Sci Tech* 2000, 19:509-543). This latter type of vaccines has the typical drawbacks associated with the use of live attenuated vaccines, specifically, the risk of mutations reverting the virulence of the virus or making it lose its immunogenicity.

Recombinant subunit vaccines containing the IBDV protein VP2 expressed in several expression systems, for example, bacteria, yeasts or baculovirus, usually in fusion protein form, have been disclosed. The results obtained in chicken immunization tests with said vaccines have not been completely satisfactory.

Empty viral capsids or virus-like particles (VLPs) constitute an alternative to the use of live attenuated vaccines and of recombinant subunit vaccines. VLPs are obtained by self-assembly of the subunits constituting the viral capsid and mimicking the structure and antigenic properties of the native virion, even though they lack genetic material, as a result of which they are incapable of replicating themselves. Apart from their application for vaccination purposes, VLPs can be used as vectors of molecules of biological interest, for example, nucleic acids, peptides or proteins. By way of illustration, parvovirus VLPs (US 6,458,362) or human immunodeficiency syndrome (HIV) VLPs (US 6,602,705), can be mentioned.

Morphogenesis is a vital process for the viral cycle requiring successive steps associated to modifications in the polypeptide precursors. As a result, viruses have developed strategies allowing the sequential and correct interaction between each one of their components. One of these strategies, frequently used by icosahedral viruses, is the use of polypeptides coming from a single polyprotein as the base of their structural components. In these cases, the suitable proteolytic processing of said polyprotein plays a crucial role in the assembly process.

This concept for the assembly of IBDV capsids has been demonstrated in earlier work (Fernández-Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles. *Journal of General Virology* 79, 1047-1054). Expression of the gene encoding for the IBDV polyprotein in eukaryotic cells gives rise to the formation of VLPs that are completely morphologically and biochemically indistinguishable from the IBDV virions. It has also been shown that the assembly of the capsids requires only the synthesis and correct processing of the viral polyprotein and is independent of the presence of the viral genome or of other proteins encoded by the viral genome, such as VP5 and VP1.

Parallely to the formation of capsids, the IBDV VP4 product is able to self-assemble in tubular structures of 20 nm in diameter. These tubules, known as type II tubules, are partially copurified with the viral particles. Other experiments have demonstrated that obtaining IBDV VLPs, using recombinant baculoviruses (rBVs) for polyprotein expression, is extremely inefficient, the accumulation of large amounts of type I tubules in the cytosol of the infected cells being obtained (Martínez-Torrecuadrada, J. L., Castón, J. R., Castro, M., Carrascosa, J. L., Rodríguez, J. F. & Casal, J. I. (2000). Different architectures in the assembly of infectious bursal disease virus capsid proteins expressed in insect cells. *Virology* 278:322-331; Chevalier, C., Lepault, J., Erk, I., Da Costa, B. & Delmas, B. (2002). The maturation process of pVP2 requires assembly of infectious bursal disease virus capsids. *Journal of Virology* 76:2384-2392). It has recently been demonstrated that this is due to the proteolytic cleavage to which the VP3 protein is subjected when it is synthesized in insect cells. Said proteolysis is almost completely prevented by the formation of VP3/VP1 complexes (Maraver, A., Oña, A., Abaitua, F., González, D., Clemente, R., Diaz-Ruiz, A., Caston, J. R., Pazos, F. & Rodríguez, J. F. (2003). The oligomerization domain of VP3, the scaffolding protein of infectious bursal disease virus, plays a critical role for capsid formation. *Journal of Virology* 77:6438-6449).

Therefore, the results obtained to date from the IBDV gene expression in different recombinant systems has allowed concluding that: (i) the assembly process is independent of the presence of genetic material of the virus, (ii) only the polypeptides encoded by the polyprotein gene are necessary for the assembly, and (iii) the assembly requires a
5 coordinated interaction between the pVP2 and VP3 peptides.

However, it is not known if the VP2/VP3 interaction is established between VP2 and VP3 domains of the polyprotein precursor when it has not yet undergone modifications, or on the contrary, if this interaction occurs after the processing of the precursor. Furthermore, current information does not exclude the possibility that VP4 could play a relevant role in
10 the morphogenesis of the viral capsid. In fact, IBDV VLPs formed by assembly of the IBDV VP2, VP3 and VP4 proteins have been disclosed (US 6,528,063, US 5,788,970 and JP 5194597).

On the other hand, information regarding IBDV protein expression by means of genetic engineering in different cell models is scarce. IBDV protein expression in insect
15 cells, bacteria and yeasts has been disclosed. Jagadish et al. (Jagadish MN, Vaughan PR, Irving RA, Azad AA, Macreadie IG. (1990). Expression and characterization of infectious bursal disease virus polyprotein in yeast. *Gene* 9:179-186; Macreadie IG, Vaughan PR, Chapman AJ, McKern NM, Jagadish MN, Heine HG, Ward CW, Fahey KJ, Azad AA. (1990). Passive protection against infectious bursal disease virus by viral VP2 expressed in
20 yeast. *Vaccine* 8:549-552) disclose IBDV VP2 expression in yeasts. The disclosed results indicate that the viral polyprotein expression in two yeast species, *Saccharomyces cerevisiae* and *Saccharomyces pombe*, is very inefficient, a large variety of protein products of different molecular mass being accumulated. The failure to obtain protein products of the expected size and the inability to detect structures produced due to the assembly of the latter was
25 attributed to two possible causes: (i) possible toxicity of the IBDV protease (VP4) in this system; and/or (ii) the inefficiency of the expression system to carry out a correct transcription and/or translation of the IBDV polyprotein. Recently, Pitcovski et al. (Pitcovski J., Gutter B, Gallili G, Goldway M, Perelman B, Gross G, Krispel S, Barbakov M, Michael A. (2003). *Vaccine* 21:4736-4743) have disclosed the IBDV VP2 expression in *Pichia*
30 *pastoris* and the immunization of chickens with a material comprising the recombinant protein (rVP2) in partially purified form. In no case has the obtainment of IBDV VLPs in yeast been disclosed.

Earlier work developed by the inventors has enabled establishing systems for obtaining IBDV VLPs using different eukaryotic expression vectors. These vectors have been used for IBDV polyprotein expression in the absence or presence of the viral VP1 RNA polymerase. The biochemical characterization of purified VLPs demonstrates that they contain pVP2, VP2 and VP3 proteins when only the viral polyprotein is expressed, and the pVP2, VP2, VP3 and VP1 proteins when the simultaneous expression of the polyprotein and viral RNA polymerase is carried out (Fernández-Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles. *Journal of General Virology* 79: 1047-1054; Martínez-Torrecuadrada, J. L., Castón, J. R., Castro, M., Carrascosa, J. L., Rodríguez, J. F. & Casal, J. I. (2000). Different architectures in the assembly of infectious bursal disease virus capsid proteins expressed in insect cells. *Virology* 278: 322-331; Maraver, A., *et al.*, (2003) cited *supra*; Lombardo, E., Maraver, A., Castón, J. R., Rivera, J., Fernández-Arias, A., Serrano, A., Carrascosa, J. L. & Rodríguez, J. F. (1999). VP1, the putative RNA-dependent RNA polymerase of infectious bursal disease virus, forms complexes with the capsid protein VP3, leading to efficient encapsidation into virus-like particles. *Journal of Virology* 73: 6973-6983). However, VLPs solely based on IBDV pVP2 and VP3, or their potential use for vaccine purposes or as vehicles of biological products of interest, have not been previously disclosed.

SUMMARY OF THE INVENTION

The invention is faced with the problem of providing new effective and safe vaccines against the infectious bursal disease virus (IBDV).

The solution provided by this invention is based on it being possible to obtain correctly assembled IBDV VLPs by means of the simultaneous expression of the IBDV pVP2 and VP3 polypeptides as independent genes and as the only representation of IBDV proteins in a gene expression system. Said VLPs are formed by self-assembly of only IBDV pVP2 and VP3, whereby they lack IBDV VP4 and, for this reason, are called VLP(-VP4) (singular) or VLPs(-VP4) (plural) in this description. Said VLPs(-VP4) can be used for example for therapeutic or diagnostic purposes, for example, in the manufacture of vaccines to protect birds from the infection caused by IBDV, or in the manufacture of gene therapy vectors.

The obtained results allow conceiving two new conclusions to the understanding of the IBDV assembly pattern: (i) the interactions between the pVP2/VP3 polypeptides result in an efficient assembly of the IBDV particles without needing expression of the whole polyprotein, and (ii) the VP4 polyprotein is not required for capsid formation.

5 These results have allowed designing a new strategy or process for the efficient production of IBDV VLPs containing antigenically relevant protein elements so as to induce an immune response. This strategy is based on the use of a gene expression system or vector allowing the coexpression of pVP2 and VP3 polypeptides as independent genes and preventing the synthesis of the polyprotein precursor of said polypeptides as well as the
10 presence of the VP4 polypeptide during the empty viral capsid assembly process.

The expression and obtainment of IBDV VLPs, particularly VLPs(-VP4), in insect cells is described in a particular embodiment, while in another particular embodiment, said VLPs(-VP4) are obtained in yeasts, with a very high yield and a very low economic cost.

The vaccines obtained using said VLPs(-VP4) have a number of advantages since, on
15 one hand, the handling of highly infectious material is prevented, the potential risk of the occurrence of new IBDV mutants is prevented and the use of a live virus in poultry farms is eliminated, thus preventing the risk of spreading IBDV vaccine strains to the environment, and on the other hand, it enables the development of differential diagnostic systems to discriminate between vaccinated and infected animals. These diagnostic systems are based
20 on the detection of antibodies against VP2 and VP4 proteins. Animals with IBDV develop a strong humoral response to both proteins. However animals immunized with VLPs(-VP4) only have antibodies against the VP2 protein.

Consequently, an object of the present invention consists of an empty IBDV viral capsid, VLP(-VP4), with immunogenic activity against infection in IBDV, characterized in
25 that it is constituted by self-assembly of only IBDV pVP2 and VP3 proteins.

A further aspect of this invention is related to a process for producing said IBDV VLPs(-VP4) provided by this invention, based on the gene coexpression of said IBDV pVP2 and VP3 proteins as two independent genes.

The nucleic acids, gene constructs, expression systems and host cells developed for
30 implementing said process of producing said IBDV VLPs(-VP4), as well as their use for the production of said IBDV VLPs(-VP4), constitute further aspects of the present invention.

Said IBDV VLPs(-VP4) have the ability to immunize animals, particularly birds, against the avian disease caused by IBDV, as well as the ability to incorporate in vectors or

vehicles molecules of biological interest, for example, polypeptides, proteins, nucleic acids, etc. In a particular embodiment, said IBDV VLPs(-VP4) can be used in the manufacture of a vaccine to protect birds against the virus causing the avian disease known as infectious bursal disease (IBDV). Virtually any bird, preferably those avian species of economic
5 interest, for example, chickens, turkeys, geese, ganders, pheasants, quails, ostriches, etc., can be immunized against the infection caused by IBDV with the vaccines provided by this invention. In another particular embodiment, said IBDV VLPs(-VP4) can internally incorporate into vehicles products with biological activity, for example, nucleic acids, peptides, proteins, drugs, etc., whereby they can be used in the manufacture of gene therapy
10 vectors.

Therefore, in a further aspect, the present invention is related to the use of said IBDV VLPs(-VP4), in the manufacture of medicaments, such as vaccines and gene therapy vectors. Said vaccines and vectors constitute further aspects of the present invention. In a particular embodiment, said vaccine is a vaccine useful for protecting birds from the infection caused
15 by IBDV. In a specific embodiment, said birds are selected from the group formed by chickens, turkeys, geese, ganders, pheasants, quails and ostriches, preferably chickens.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1. (a) The diagram schematizes the proteolytic processing steps necessary for
20 the formation of mature VP2 and VP3 capsid proteins from the polyprotein precursor. (b) The diagram reflects the different genetic constructs derived from the IBDV polyprotein described up until now, as well as the structures produced by means of its expression in different heterologous systems. The numbers indicate the position corresponding to the first and last amino acid residue of the polyprotein present in each one of the constructs. The
25 lower portion of the figure shows images obtained by means of transmission electron microscopy of the structures obtained by means of expression of the different constructs. The bar corresponds to 50 nm. The data has been taken from the following literature references: Fernández-Arias *et al.*, (1998), cited *supra*; Maraver *et al.*, (2003), cited *supra*; Martínez-Torrecuadrada *et al.*, (2000), cited *supra*; Castón *et al.*, 2001. C terminus of infectious bursal
30 disease virus major capsid protein VP2 is involved in definition of the t number for capsid assembly. *Journal of Virology* 75, 10815-10828.

Figure 2. Microscopic analysis of H5 insect cells coexpressing pVP2 and VP3. The pVP2 and VP3 protein subcellular distribution was analyzed by means of confocal

immunomicroscopy. Cells infected with the FB/pVP2 (a), FB/VP3 (b), or FBD/pVP2-VP3 (c-e) rBVs were incubated with anti-pVP2 rabbit serum and anti-VP3 rat serum. Then the cells were incubated with goat anti-rabbit IgG serum coupled to Alexa 488 (red) and goat anti-rat IgG serum coupled to Alexa 594 (green). The cores were stained with To-Pro 3 (blue). (e) Overlaying of the images shown in panels (c) and (d). Electron microscopy images corresponding to sections of H5 cells infected with different genetic constructs derived from the IBDV polyprotein. (f) Low-magnification image of an H5 cell infected with a parental Fb virus. The insert corresponds to an enlarged detail of the area indicated by the box. (g) Low-magnification image of an H5 cell infected with the FBD/pVP2-VP3 virus. The insert corresponds to an enlarged detail of the area indicated by the box. (h) High-magnification image of an H5 cell infected with the FBD/pVP2-VP3 virus showing the formation of IBDV structures in detail. (i) High-magnification image of a BSC1 cell infected with the VTLacOI/POLY recombinant vaccine virus showing structures similar to those detected in panel (h). The bars indicate 600 nm (panels f and g) and 200 nm (panels h and i).

Figure 3. Structural and biochemical characterization of the structures derived from IBDV produced in insect cells coinfecting with the (rBV) FB/pVP2 + FB/his-VP3 recombinant baculoviruses. Cells coinfecting with FB/pVP2 and FB/his-VP3 rBVs, or infected with the FBD/Poly-VP1 or FB/pVP2 virus, were used to purify structures derived from IBDV by means of centrifugation on sucrose gradients. Panels (a), (b), and (c) show transmission electron microscopy images corresponding to fraction 4 of the gradients obtained from infections with FBD/Poly-VP1, FB/pVP2+FB/his-VP3, and FB/pVP2, respectively. Panel (d) shows the results of a Western blot analysis of the sucrose gradients corresponding to the cultures infected with FBD/Poly-VP1 and FB/pVP2+FB/his-VP3, respectively. The total extracts (input) and the different fractions of the sucrose gradients (fraction F1 corresponds to the bottom of the gradient) were analyzed by means of Western blot using specific sera against the IBDV VP1, pVP2, VP3, and VP4 proteins, respectively. The molecular mass of the immunoreactive polypeptides is indicated in kDa.

Figure 4. Biochemical and structural characterization of IBDV VLPs produced in *S. cerevisiae* transformed with the plasmid pESCURA/pVP2-VP3-GFP. A *S. cerevisiae* culture transformed with the plasmid pESCURA/pVP2-VP3-GFP was grown at 30°C in a medium supplemented with the inducer galactose. At 18 hours, the culture was harvested and centrifuged. The resulting sediment was processed by means of fractioning in a 25-50% linear sucrose gradient. A) Biochemical analysis of samples corresponding to the

sediment before fractioning (T) as well as the different fractions of the sucrose gradient. The samples were analyzed by means of SDS-PAGE and Western blot using specific antibodies against VP3 (anti-VP3) and pVP2 (anti-pVP2) proteins. The arrows indicate the positions of the immunoreactive bands corresponding to the VP3-GFP (61 kDa) and pVP2 (48 kDa) proteins, respectively. B) The structural analysis of the obtained samples was carried out by means of TEM. The image corresponds to a micrograph obtained from an aliquot corresponding to the mixture of fractions 7, 8 and 9 of the sucrose gradient. The sample was stained with uranyl acetate and observed by means of TEM. The bar corresponds to 65 nm. C) VLPs sample obtained by means of the IBDV polyprotein expression in mammal cells by means of infection with the VT7/Poly recombinant vaccine virus (Fernández-Arias *et al.*, (1998), cited *supra*). The bar corresponds to 65 nm.

DETAILED DESCRIPTION OF THE INVENTION

In a first aspect, the invention provides an empty capsid of the infectious bursal disease virus (IBDV), hereinafter VLP(-VP4) of the invention, characterized in that it is constituted by assembly of only IBDV pVP2 proteins and IBDV VP3 proteins.

The term "IBDV", as it is used in the present invention, refers to the different IBDV strains belonging to any of the known serotypes (1 or 2) [by way of illustration, see the review carried out by van den Berg TP, Etteradossi N, Toquin D, Meulemans G., en *Rev Sci Tech* 2000 19: 509-43] and the terms "IBDV pVP2 protein" and "IBDV VP3 protein" refer to the different forms of the pVP2 and VP3 proteins representative of any of the mentioned IBDV strains [NCBI protein databank], according to the definition made by Sánchez and Rodríguez (1999) (Sánchez AB, Rodríguez JF. Proteolytic processing in infectious bursal disease virus: identification of the polyprotein cleavage sites by site-directed mutagenesis. *Virology*. 1999 Sep 15; 262(1):190-199), as well as proteins substantially homologous to said IBDV pVP2 and VP3 proteins, i.e. proteins the amino acid sequences of which have a degree of identity regarding said IBDV pVP2 and VP3 proteins of at least 60%, preferably of at least 80%, more preferably of at least 90% and even more preferably of at least 95%.

The IBDV pVP2 protein present in the VLP(-VP4) of the invention can be any pVP2 protein representative of any IBDV strain, for example, the full-length pVP2 protein of IBDV Soroa strain [NCBI, access number AAD30136]

The IBDV VP3 protein present in the VLP(-VP4) of the invention can be any VP3 protein representative of any IBDV strain, for example, the full-length VP3 protein of IBDV Soroa strain [NCBI, access number AAD30136].

In a particular embodiment, the VLPs(-VP4) of the invention have a diameter of 65-
5 70 nm and a polygonal contour indistinguishable from the IBDV VLPs obtained in other expression systems (Figure 4C).

The VLPs(-VP4) of the invention can be obtained by means of the simultaneous expression of said IBDV pVP2 and VP3 proteins in suitable host cells. Said suitable host cells are cells containing the nucleotide sequence encoding the IBDV pVP2 protein and the
10 nucleotide sequence encoding the IBDV VP3 protein, either in a single gene construct or in two gene constructs. In a particular embodiment, said suitable host cells are cells that are transformed, transfected or infected with a suitable expression system, such as an expression system comprising a gene construct, wherein said gene construct comprises the nucleotide sequence encoding for the IBDV pVP2 protein and the nucleotide sequence encoding the
15 IBDV VP3 protein, or else alternatively with an expression system comprising a first gene construct comprising the nucleotide sequence encoding for the IBDV pVP2 protein, and a second gene construct comprising the nucleotide sequence encoding for the IBDV VP3 protein.

Therefore, in another aspect, the invention provides a nucleic acid, the nucleotide
20 sequence of which comprises the nucleotide sequence encoding for said IBDV pVP2 protein and the nucleotide sequence encoding for said IBDV VP3 protein in the form of two independent genes. More specifically, the nucleic acid provided by this invention is characterized in that its nucleotide sequence is constituted by (i) a nucleotide sequence comprising the open reading frame corresponding to the IBDV pVP2 protein and (ii) a
25 nucleotide sequence comprising the open reading frame corresponding to the IBDV VP3 protein. One feature of the nucleic acid provided by this invention is that it lacks the nucleotide sequence comprising the open reading frame corresponding to the IBDV VP4 protein. As it is used in this description, the term "open reading frame corresponding to the pVP2 protein" or "open reading frame corresponding to the IBDV VP3 proteins" includes,
30 apart from the nucleotide sequences of said open reading frames, other open reading frames analogous to the same encoding frames of the IBDV pVP2 and VP3 proteins. As it is used herein, the term "analogous" intends to include any nucleotide sequence which can be isolated or constructed on the base of the encoding nucleotide sequence of IBDV pVP2 and

VP3, for example by means of the introduction of conservative or non-conservative nucleotide replacements, including the insertion of one or more nucleotides, the addition of one or more nucleotides at any of the ends of the molecule, or the deletion of one or more nucleotides at any end or inside of the sequence. Generally, a nucleotide sequence analogous to another nucleotide sequence is substantially homologous to said nucleotide sequence. In the sense used in this description, the expression "substantially homologous" means that at the nucleotide level the nucleotide sequences in question have a degree of identity of at least 60%, preferably of at least 80%, more preferably of at least 90%, and even more preferably of at least 95%.

10 In another aspect, the invention provides a gene construct comprising a nucleic acid provided by this invention, i.e. a nucleic acid the nucleotide sequence of which is constituted by (i) a nucleotide sequence comprising the open reading frame corresponding to the IBDV pVP2 protein and (ii) a nucleotide sequence comprising the open reading frame corresponding to the IBDV VP3 protein. Said gene construct lacks the nucleotide sequence comprising the open reading frame corresponding to the IBDV VP4 protein.

15 In another aspect, the invention provides an expression vector or system selected from:

- 20 a) an expression system comprising a gene construct provided by this invention, operatively bound to transcription, and optionally translation, control elements, wherein said gene construct comprises the nucleotide sequence comprising the open reading frame corresponding to the IBDV pVP2 protein and the nucleotide sequence comprising the open reading frame corresponding to the IBDV VP3 protein; and
- 25 b) an expression system comprising two gene constructs, a first gene construct comprising the open reading frame corresponding to the IBDV pVP2 protein, operatively bound to transcription, and optionally translation, control elements, and a second gene construct comprising the open reading frame corresponding to the IBDV VP3 protein, operatively bound to transcription, and optionally
- 30 translation, control elements.

In a particular embodiment, the expression system provided by this invention comprises a gene construct comprising (i) a nucleotide sequence comprising the open reading frame or encoding region corresponding to the IBDV pVP2 protein and (ii) a nucleotide sequence comprising the open reading frame or encoding region corresponding to the IBDV VP3 protein, wherein said gene construct is operatively bound to transcription, and optionally translation, control elements.

In another particular embodiment, the expression system provided by this invention comprises (i) a first gene construct, operatively bound to transcription, and optionally translation, control elements, said first gene construct comprising a nucleotide sequence comprising the open reading frame or encoding region corresponding to the IBDV pVP2 protein, and (ii) a second gene construct, operatively bound to transcription, and optionally translation, control elements, said second gene construct comprising a nucleotide sequence comprising the open reading frame or encoding region corresponding to the IBDV VP3 protein.

The transcription, and optionally translation, control elements present in the expression system provided by this invention include promoters, directing the transcription of the nucleotide sequences of interest to which they are operatively linked, and other sequences necessary or suitable for the transcription and its suitable regulation in time and place, for example, start and end signals, cleavage sites, polyadenylation signal, replication origin, transcriptional activators (enhancers), transcriptional silencers (silencers), etc.

Virtually any suitable expression system or vector can be used in the generation of the expression system provided by this invention. By way of illustration, said suitable expression or vector systems can be selected, according to the conditions and needs of each specific case, from plasmids, bacmids, yeast artificial chromosomes (YACs), bacteria artificial chromosomes (BACs), bacteriophage P1-based artificial chromosomes (PACs), cosmids, or viruses, which can further have a heterologous replication origin, for example, bacterial or of yeast, so that it may be amplified in bacteria or yeasts, as well as a marker usable for selecting the transfected cells different from the gene or genes of interest. These expression systems or vectors can be obtained by conventional methods known by persons skilled in the art [Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory] and form part of the present invention. In a particular embodiment, said expression or vector system is a plasmid, such as a plasmid suitable for transforming yeasts, for example, the plasmid called

pESCURA/pVP2-VP3-GFP (Example 2), or a virus, such as a recombinant baculovirus (rBV), for example, the rBV called FBD/pVP2-his-VP3 (Example 1.2), simultaneously expressing both proteins (IBDV pVP2 and his-VP3) in insect cells during the replication cycle, or the rBVs called FB/pVP2 and FB/his-VP3 (Example 1.1) expressing the IBDV
5 pVP2 and his-VP3 proteins, respectively, when coinfecting insect cells, IBDV VLPs(-VP4) being obtained.

In another aspect, the invention provides a host cell containing the encoding nucleotide sequence of the IBDV pVP2 protein and the encoding nucleotide sequence of the IBDV VP3 protein, either in a single gene construct or in two different gene constructs. In a
10 particular embodiment, said host cell is a host cell that is transformed, transfected or infected with (i) an expression system provided by this invention comprising either a gene construct wherein said gene construct comprises the nucleotide sequence encoding for said IBDV pVP2 protein and the nucleotide sequence encoding for said IBDV VP3 protein, or else alternatively with (ii) an expression system comprising a gene construct comprising the
15 nucleotide sequence encoding for said IBDV pVP2 protein and another gene construct comprising the nucleotide sequence encoding for said IBDV VP3 protein.

In a particular embodiment, the host cell provided by this invention is a host cell that is transformed, transfected or infected with an expression system comprising a gene construct comprising (i) a nucleotide sequence comprising the open reading frame or
20 encoding region corresponding to the IBDV pVP2 protein and (ii) a nucleotide sequence comprising the open reading frame or encoding region corresponding to the IBDV VP3 protein, wherein said gene construct is operatively bound to transcription, and optionally translation, control elements.

In another particular embodiment, the host cell provided by this invention is a host
25 cell that is transformed, transfected or infected with a first gene construct, operatively bound to transcription, and optionally translation, control elements, said first gene construct comprising a nucleotide sequence comprising the open reading frame or encoding region corresponding to the IBDV pVP2 protein, and with a second gene construct, operatively bound to transcription, and optionally translation, control elements, said second gene
30 construct comprising a nucleotide sequence comprising the open reading frame or encoding region corresponding to the IBDV VP3 protein.

Virtually any host cell susceptible to being transformed, transfected or infected by an expression system provided by this invention can be used, for example, mammal cells, bird

cells, insect cells, yeasts, etc; however, in a particular embodiment, said host cell is selected from yeasts and insect cells. Yeasts are suitable due to the simplicity and production cost. Insect cells are suitable when the expression system comprises one or two recombinant baculoviruses (rBV). The use of rBV is advantageous due to biosafety issues related to the host
5 range of the baculoviruses, incapable of replicating in other cell types which are not insect cells.

In a particular embodiment, the invention provides a host cell, such as a yeast, for example, *Saccharomyces cerevisiae*, *Saccharomyces pombe*, etc., transformed with an expression system, such as a plasmid or an expression vector, comprising a gene construct
10 provided by this invention comprising the nucleotide sequence encoding for said IBDV pVP2 protein and the nucleotide sequence encoding for the IBDV VP3 protein.

In another particular embodiment, the invention provides a host cell, such as an insect cell, infected with an expression system, such as a recombinant baculovirus, comprising a gene construct provided by this invention comprising the nucleotide sequence encoding for said
15 IBDV pVP2 protein and the nucleotide sequence encoding for the IBDV VP3 protein.

In another particular embodiment, the invention provides a host cell, such as an insect cell, coinfecting with an expression system comprising a first recombinant baculovirus comprising a gene construct provided by this invention comprising the nucleotide sequence encoding for said IBDV pVP2 protein and with a second recombinant baculovirus comprising
20 a gene construct provided by this invention comprising the nucleotide sequence encoding for the IBDV VP3 protein.

In another aspect, the invention provides a process for the production of VLPs(-VP4) of the invention comprising culturing a host cell provided by this invention containing the encoding nucleotide sequence of said IBDV pVP2 protein and the encoding nucleotide
25 sequence of said IBDV VP3 protein, either in a single gene construct or in two different gene constructs, and simultaneously expressing said IBDV pVP2 and VP3 proteins, and if so desired, recovering said VLPs(-VP4) of the invention. In a particular embodiment, said host cell provided by this invention is a cell that is transformed, transfected or infected with a suitable expression system, such as an expression system comprising a gene construct, wherein
30 said gene construct comprises the nucleotide sequence encoding for said IBDV pVP2 protein and the nucleotide sequence encoding for said IBDV VP3 protein, or else alternatively with an expression system comprising a first gene construct comprising the nucleotide sequence

encoding for said IBDV pVP2 protein and a second gene construct comprising the nucleotide sequence encoding for said IBDV VP3 protein.

Said process therefore comprises the gene coexpression of said IBDV pVP2 and VP3 proteins as two independent genes. After the simultaneous expression of said proteins (IBDV pVP2 and VP3) in said cells, the expressed proteins are assembled and form the VLPs(-VP4) of the invention, which can be isolated or withdrawn from the medium and purified if desired. The isolation and purification of said VLPs(-VP4) of the invention can be carried out by means of conventional methods, for example, by means of fractioning on sucrose gradients.

In a particular embodiment, the simultaneous gene coexpression of IBDV pVP2 and VP3 proteins is carried out by means of the use of an rBV allowing the simultaneous expression of said proteins from two independent chimeric genes in insect cells. In this case, the production of VLPs(-VP4) of the invention can be carried out by means of a process comprising, first, the obtainment of a gene expression system made up of an rBV containing a gene construct simultaneously encoding for said IBDV pVP2 and VP3 proteins, such as the rBV called FBD/pVP2-his-VP3 (Example 1.2), or alternatively the obtainment of an rBV containing a gene construct encoding for the IBDV pVP2 protein and the obtainment of another rBV containing a gene construct encoding for said IBDV VP3 protein, such as the rBVs called FB/pVP2 and FB/his-VP3 (Example 1.1), followed by the infection of insect cells with said expression system based on said recombinant baculovirus(es), expression of the recombinant proteins and if so desired, isolation of the VLPs(-VP4) of the invention formed by assembly of said IBDV pVP2 and VP3 proteins, and optionally subsequent purification of said VLPs(-VP4) of the invention.

The construction of a recombinant baculovirus allowing the independent expression of the IBDV pVP2 and VP3 proteins can be carried out by any person skilled in the art based on that described herein and on the state of the art concerning this technology (Cold Spring Harbor, N.Y.; Leusch MS, Lee SC, Olins PO. 1995. A novel host-vector system for direct selection of recombinant baculoviruses (bacmids) in *Escherichia coli*. Gene 160: 191-4; Luckow VA, Lee SC, Barry GF, Olins PO. 1993. Efficient generation of infectious recombinant baculoviruses by site-specific transposon-mediated insertion of foreign genes into a baculovirus genome propagated in *Escherichia coli*. J Virol 67: 4566-79).

In another particular embodiment, the gene coexpression of the IBDV pVP2 and VP3 proteins is carried out by means of a vector allowing the expression of said proteins in yeast cells. In this case, the production of VLPs(-VP4) of the invention can be carried out by means

of a process comprising, first, the obtainment of a gene expression system made up of a plasmid containing a gene construct simultaneously encoding for the IBDV pVP2 and VP3 proteins, followed by the transformation of yeasts with said expression system, expression of the recombinant proteins, and if so desired, isolation of the VLPs(-VP4) of the invention
5 formed by assembly of said IBDV pVP2 and VP3 proteins, and optionally subsequent purification of said VLPs(-VP4) of the invention. In a specific embodiment, the suitable expression system for transforming yeasts is based on a pESC Yeast (Stratagene) expression system such as, for example, the plasmid pESCURA/pVP2/VP3-GFP (Example 2) containing a gene construct encoding for the IBDV pVP2 and VP3-GFP proteins.

10 The obtainment of yeasts transformed with a gene construct or with a suitable expression system or vector allowing the simultaneous expression of the IBDV pVP2 and VP3 proteins can be carried out by any person skilled in the art based on that described herein and on the state of the art concerning this technology (pESC epitope tagging vectors Instructions manual. Stratagene www.stratagene.com; Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989).
15 Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory).

In another aspect, the invention is related to the use of the gene expression system provided by this invention for producing and obtaining the VLPs(-VP4) of the invention.

The VLPs(-VP4) of the invention can be used to immunize animals, particularly birds, *per se* or as vectors or vehicles of molecules with biological activity, for example,
20 polypeptides, proteins, nucleic acids, drugs, etc., whereby they can be used for therapeutic or diagnostic purposes. In a particular embodiment, said molecules with biological activity include antigens or immune response inducers in animals or humans to whom they are supplied, or drugs which can be released in their specific action site, or nucleic acid sequences, all being useful in gene therapy and intended for being introduced inside the suitable cells.

25 Therefore, in another aspect, the invention is related to the use of the VLPs(-VP4) of the invention in the manufacture of medicaments such as vaccines, gene therapy vectors (delivery systems), etc. In a particular embodiment, said medicament is a vaccine intended for conferring protection to animals, particularly birds, against the infectious bursal disease virus (IBDV). In another particular embodiment, said medicament is a gene therapy vector.

30 In another aspect, the invention provides a vaccine comprising a therapeutically effective amount of VLPs(-VP4) of the invention, optionally together with one or more pharmaceutically acceptable adjuvants and/or vehicles. Said vaccine is useful for protecting animals, particularly birds, against the infectious bursal disease virus (IBDV). In a particular

embodiment, said birds are selected from the group formed by chickens, turkeys, geese, ganders, pheasants, quails and ostriches. In a preferred embodiment, the vaccine provided by this invention is a vaccine useful for protecting chickens from the infection caused by IBDV.

In the sense used in this description, the expression "therapeutically effective amount" refers to the amount of VLPs(-VP4) of the invention calculated for producing the desired effect and will generally be determined, among others, by the characteristics of the VLPs(-VP4) and the immunization effect to be achieved.

The pharmaceutically acceptable adjuvants and vehicles which can be used in said vaccines are those adjuvants and vehicles known by the persons skilled in the art and normally used in the manufacture of vaccines.

In a particular embodiment, said vaccine is prepared in form of an aqueous solution or suspension in a pharmaceutically acceptable diluent, such as saline solution, phosphate-buffered saline solution (PBS), or any other pharmaceutically acceptable diluent.

The vaccine provided by this invention can be administered by any suitable administration route which results in a protective immune response against the heterologous sequence or epitope used, to which end said vaccine will be formulated in the dosage form suited to the chosen administration route. In a particular embodiment, the administration of the vaccine provided by this invention is carried out parenterally, for example, intraperitoneally, subcutaneously, etc.

The following Examples illustrate the invention and should not be considered limiting of the scope thereof.

EXAMPLE 1

Obtaining VLPs(-VP4) by means of the coexpression of pVP2 (pVPX) and VP3 in insect cells

1.1 Obtaining VLPs(-VP4) by means of the coexpression of pVP2 (pVPX) and VP3 with two independent rBVs in insect cells

The results of a series of experiments designed to analyze the possibility of obtaining IBDV VLPs from a strategy avoiding the synthesis of the IBDV polyprotein and, therefore, the presence of the VP4 protease during the assembly process are described in this example. The experimental design is based on the coexpression of the IBDV pVP2 and VP3 proteins from two independent chimeric genes. To that end, two recombinant baculoviruses (rBVs) described above, FB/his-VP3 (Kochan, G., González, D. & Rodríguez, J. F. (2003).

Characterization of the RNA binding activity of VP3, a major structural protein of IBDV. *Archives of Virology* 148, 723-744) and FB/VPX [also identified as FB/pVP2 in this description] (Martínez-Torrecuadrada, J. L., Castón, J. R., Castro, M., Carrascosa, J. L., Rodríguez, J. F. & Casal, J. I. (2000). Different architectures in the assembly of infectious
5 bursal disease virus capsid proteins expressed in insect cells. *Virology* 278, 322-331) have been used. These rBVs were generated by means of the cloning into suitable vectors of the complementary DNA (cDNA) encoders of the IBDV pVP2 and pVP3 proteins. Said cDNAs were obtained by RT-PCR from the A segment of the serotype I IBDV Soroa strain genome a (NCBI access number AAD30136). The rBV FB/his-VP3 expresses a chimeric VP3
10 protein which at its N-terminal end contains a tandem of six histidines fused to the VP3 sequence (Met754-Glu1012 of the polyprotein) called his-VP3. rBV FB/pVP2 expresses the encoding region of the pVP2 protein (Met1-Ala512).

The analysis of the expression of these pVP2 and his-pVP3 proteins, whether independently or together, was carried out in cell cultures. To carry out these experiments,
15 single layer cell cultures from the insect *Trichoplusia ni* (H5, Invitrogen) were used, which were grown on cover glasses. Said cultures were independently infected with FB/pVP2, FB/his-VP3, or coinfecting with both rBVs. The multiplicity of infection was 5 plaque forming units (pfu)/cell. The cells were fixed at 48 hours post-infection (h.p.i), and incubated with rabbit anti-VP2 polyclonal serum and with rat anti-VP3 polyclonal serum (Fernández-
20 Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles. *Journal of General Virology* 79, 1047-1054). After successive washings, the cover glasses were incubated with goat anti-rat serum conjugated with Alexa 594 and with goat anti-rabbit serum conjugated with Alexa 488 (Jackson Immunoresearch Laboratories, Inc.). The cellular
25 cores were stained with the specific To-Pro-3 marker (Molecular Probes, Inc.). The samples were finally viewed by epifluorescence with a Zeiss Axiovert 200 microscope equipped with the Bio Rad Radiance 2100 confocal system. The images obtained were stored using the Laser Sharp Package (Bio Rad) software equipment. As is shown in Figure 2a, in the cultures infected with FB/pVP2, the anti-VP2 serum showed a fine granular signal mixed
30 with tubular structures, both distributed throughout the cytoplasm. The anti-VP3 signal, detected in the cells infected with rBV FB/his-VP3, was characterized by the presence of spherical-shaped, and apparently hollow, accumulations around the core. In the cultures coinfecting with both rBVs, a notable modification in the distribution pattern of both proteins

was detected. In these cells, the specific signals of pVP2 and VP3 were collocated in spherical and dense accumulations, suggesting that their coexpression allowed the formation of pVP2/his-VP3 complexes (Figure 2c to 2e).

For the purpose of characterizing these structures in greater detail, similar extracts
5 corresponding to cells infected with FB/pVP2+FB/hisVP3 were analyzed by transmission
electron microscopy (TEM). As a control, and in parallel, H5 cell cultures infected with the
wild strain of the FBD (FastBacDual, Invitrogen) virus were analyzed by the same
technique. After the infection, the cells were harvested after 48 hours, and processed as has
been previously described (Fernández-Arias A, Risco C, Martínez S, Albar JP & Rodríguez
10 JF. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation
of virus-like particles. *Journal of General Virology* 79:1047-1054) for their analysis in
ultrathin sections by TRANSMISSION ELECTRON MICROSCOPY. As is shown in Figure
2, the cytoplasm of the coinfecting cells contains aggregates formed by a mixture of tubules
and structures similar to capsids (Figure 2g, 2h and 2i). These aggregates were not observed
15 in any case in the samples corresponding to cells infected with wild FBD virus (Figure 2f).
The appearance and size of the tubules, as well as of the structures similar to capsids, was
similar to those previously described in cell cultures infected with VT7/Poly, a recombinant
of the vaccinia virus expressing the gene of the IBDV polyprotein (Fernández-Arias A,
Risco C, Martínez S, Albar JP & Rodríguez JF. (1998). Expression of ORF A1 of infectious
20 bursal disease virus results in the formation of virus-like particles. *Journal of General
Virology* 79:1047-1054).

To unmistakably establish that the coexpression of pVP2 and his-VP3 enabled the
assembly and, therefore, the obtainment of VLPs-(VP4), the decision was made to purify the
formed particles. To that end, H5 cell cultures were infected with FB/pVP2+FB/his-VP3. At
25 60 h.p.i., the cells were homogenized and the extracts were separated on sucrose gradients as
previously described (Lombardo E, Maraver A, Castón JR, Rivera J, Fernández-Arias A,
Serrano A, Carrascosa JL & Rodríguez JF. (1999). VP1, the putative RNA-dependent RNA
polymerase of infectious bursal disease virus, forms complexes with the capsid protein VP3,
leading to efficient encapsidation into virus-like particles. *Journal of Virology* 73:6973-
30 6983). After their centrifugation, the gradients were fractioned, and the different fractions
were analyzed by TEM as previously described (Lombardo and *et al.*, cited *supra*). As a
control, and subject to the same process, gradients corresponding to cell extracts infected
with rBV FB/VPX or with rBV FBD/Poly-VP1, were fractioned. The recombinant virus

FBD/Poly-VP1 simultaneously expresses the VP1 polypeptide and polyprotein. As was predictable, the infection with FBD/Poly-VP1 had a result of an efficient production of VLPs (Maraver A, Oña A, Abaitua F, González D, Clemente R, Diaz-Ruiz A, Castón JR, Pazos F & Rodríguez JF. (2003). The oligomerization domain of VP3, the scaffolding protein of infectious bursal disease virus, plays a critical role for capsid formation. *Journal of Virology* 77:6438-49). On the other hand, the fractions corresponding to the cells infected with FB/VPX only contain tubules of a twisted appearance. The gradients corresponding to cells coinfecting with the rBVs FB/pVP2+FB/his-VP3 contain rigid type I tubules in the fractions near the bottom of the gradient, and VLPs-(VP4) in the central and top fractions (Figure 3b). The VLPs-(VP4) isolated from the cells coinfecting with rBV FB/pVP2+FB/his-VP3 had a diameter of 65-70 nm, as well as a typical polygonal contour, absolutely indistinguishable from the purified VLPs of cultures infected with FBD/Poly-VP1 (Maraver, A., Oña, A., Abaitua, F., González, D., Clemente, R., Diaz-Ruiz, A., Caston, J. R., Pazos, F. & Rodríguez, J. F. (2003). The oligomerization domain of VP3, the scaffolding protein of infectious bursal disease virus, plays a critical role for capsid formation. *Journal of Virology* 77:6438-49) or of the cultures infected with VT7/Poly (Fernández-Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles. *Journal of General Virology* 79, 1047-1054).

For the purpose of achieving a biochemical characterization of the obtained material, Western blot experiments were carried out in which the different fractions were compared with specific sera against the VP1, pVP2, VP3 and VP4 proteins (Fernández-Arias *et al.* 1998, Lombardo *et al.*, 2000). Cell extracts infected with IBDV were used as a control. The obtained results are shown in Figure 3d. As was expected, the bands corresponding to the VP1 and VP4 polypeptides were only detected in samples corresponding to cells infected with FBD/Poly-VP1. The patterns corresponding to pVP2/VP3 in samples corresponding to cells infected with FBD/Poly-VP1 or coinfecting with FB/VPX+ FB/his-VP3 were similar, two bands corresponding to pVP2 and VP3, respectively, being detected.

1.2 Obtaining VLPs(-VP4) by means of the coexpression of pVP2 (pVPX) and VP3 with an rBV in insect cells

Furthermore, the construction of the plasmid pFBD/pVP2-his-VP3 was carried out. The first step of the construction was carried out by means of the cloning of the encoding

region of the pVP2 protein into the pFBDual vector (Invitrogen). The DNA fragment corresponding to pVP2 was obtained by means of PCR with the oligonucleotides identified as Oligo I (SEQ ID NO: 1) and Oligo II (SEQ ID NO: 2) using the plasmid pVOTE.2/Poly as a mold (Fernández-Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus infectious results in the formation of virus-like particles. *Journal of General Virology* 79, 1047-1054). The fragment was purified, subjected to digestion with the BglII and HindIII enzymes and cloned into the pFBDual vector (Invitrogen) previously digested with the BamHI and Hindi enzymes. The resulting plasmid was called pFBD/pVP2. Then, a DNA fragment containing the open reading frame corresponding to the VP3 protein was obtained by means of digestion of the plasmid pFB/his-VP3 (Kochan et al., 2003, cited *supra*) with the RsrII enzyme, treatment with Klenow, and subsequent restriction with KpnI. This DNA fragment was purified and cloned into the plasmid pFBD/pVP2 previously digested with the SmaI and KpnI enzymes. The resulting plasmid was called pFBD/pVP2-his-VP3 (SEQ ID NO: 3) and contains the encoding nucleotide sequence of the pVP2 and his-pVP3 proteins (the latter is encoded by the complementary chain to the nucleotides 6734-7585 of SEQ ID NO: 3). The amino acid sequence of the pVP2 protein and of the his-VP3 fusion protein (pVP2-his-VP3) encoded by the nucleotide sequence contained in said plasmid pFBD/pVP2-his-VP3 is shown in SEQ ID NO: 4.

The plasmid pFBD/pVP2-his-VP3 allows obtaining an rBV, called FBD/pVP2-his-VP3, expressing both proteins simultaneously during its replication cycle [<http://invitrogen.com/content/sfs/manuals/bevtest.pdf>].

The results obtained with FBD/pVP2-his-VP3 are identical to those obtained by means of the coinfection with rBVs FB/pVP2 and FD/his-VP3, IBDV VLPs(-VP4) being obtained.

EXAMPLE 2

Obtaining VLPs(-VP4) by means of the coexpression of pVP2 and VP3 as two independent genes in yeasts

For the purpose of studying the possibility of obtaining IBDV VLPs(-VP4) in yeast cultures (*Saccharomyces cerevisiae*) the vector pESCURA/pVP2-VP3-GFP was generated with the heterologous GFP gene bound to the VP3 N-terminal end. The first step in the construction of the vector was carried out by means of the cloning of the encoding region of

the IBDV pVP2 protein into the vector pESCURAinv. The plasmid pESCURAinv was generated by means of digestion of the vector pRS426 (Stratagene) with the PvuII enzyme and religation of the digestion mixture. The resulting vector, pESCURAinv, contains the multiple cloning region in reversed position with regard to that of parent vector pRS426. The

5 DNA fragment corresponding to the pVP2 protein was obtained by means of PCR with the oligonucleotides called Oligo III (SEQ ID NO: 5) and Oligo IV (SEQ ID NO: 6) using the plasmid pVOTE.2/Poly as a mold (Fernández-Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles. *Journal of General Virology* 79, 1047-1054). The

10 fragment was purified subjected to digestion with the BglII and HindIII enzymes and cloned into the vector pESCURA.inv, previously digested with the BamHI and HindIII enzymes. The resulting plasmid was called pESCURA/pVP2.

The plasmid pFB/VP3-GFP was constructed in two stages. The first one consisted of the cloning of a DNA fragment, generated by means of PCR, containing the ORF of the

15 IBDV VP3 protein lacking the termination codon. This PCR was carried out using the oligonucleotides called Oligo V (SEQ ID NO: 7) and Oligo VI (SEQ ID NO: 8) and using the plasmid pVOTE.2/Poly as a mold (Fernández-Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles. *Journal of General Virology* 79, 1047-1054). The

20 resulting DNA was digested with the EcoRI and BamHI enzymes and cloned into the vector pEGFP-N3 (Clontech), also digested with the same enzymes. The resulting plasmid was called pVP3-GFP. Then, the plasmid pEGFP-GFP was digested with the EcoRI and NotI enzymes and cloned into the vector pFastBac1 (Invitrogen). The resulting plasmid was called pFB/VP3-GFP.

25 Then, a DNA fragment that contained the open reading frame corresponding to the IBDV VP3 protein fused to the encoding region of the EGFP protein was obtained by means of digestion of the plasmid pFB/VP3-GFP with the EcoRI and NotI enzymes. This DNA fragment was purified and cloned into the plasmid pESCURA/pVP2 previously digested with the EcoRI and NotI enzymes. The resulting plasmid was called pESCURA/pVP2-VP3-

30 GFP (SEQ ID NO: 9) and contains the ORFs of the pVP2 and VP3-GFP proteins under the transcriptional control of two independent promoters, GAL 1 and GAL 10, both inducible by galactose (the pVP2 protein is encoded by the chain of nucleotides complementary to the nucleotides 5862-7343 of SEQ ID NO: 9). The amino acid sequence of the pVP2 protein and

of the VP3-GFP fusion protein (pVP2-VP3-GFP) encoded by the nucleotide sequence contained in said plasmid pESCURA/pVP2-VP3-GFP is shown in SEQ ID NO: 10.

pESCURA/pVP2-VP3-GFP was subsequently used to transform a culture of *S. cerevisiae* yeast haploid strain 499 according to a previously described protocol (Gietz, R.D. and R.A. Woods. (2002), Transformation of yeast by the Liac/SS carrier DNA/PEG method. Methods in Enzymology 350:87-96). The yeasts transformed with the plasmid were selected by means of growth on SC medium plates (CSM + YNB, 2% glucose and bacto agar) supplemented with the amino acids tryptophan, leucine and histidine and lacking uracyl (-Ura). After an incubation of 48 hours at 30°C, a colony was chosen which was used to carry out the following protein expression and VLPs-(VP4) formation analyses.

The pVP2 and VP3 protein expression and VLPs-(VP4) formation analyses were carried out following a protocol previously described for the characterization of IBDV VLPs in other expression systems (Fernández-Arias, A., Risco, C., Martínez, S., Albar, J. P. & Rodríguez, J. F. (1998). Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles. *Journal of General Virology* 79, 1047-1054; Lombardo, E., Maraver, A., Castón, J. R., Rivera, J., Fernández-Arias, A., Serrano, A., Carrascosa, J. L. & Rodríguez, J. F. (1999). VP1, the putative RNA-dependent RNA polymerase of infectious bursal disease virus, forms complexes with the capsid protein VP3, leading to efficient encapsidation into virus-like particles. *Journal of Virology* 73, 6973-698). The colony selected was cultured in liquid CSM (-Ura) + YNB medium supplemented with 2% raffinose. The culture was incubated at 30°C for 24 hours. This culture was used to inoculate, at an optical density (O.D.) of 0.2, a flask of 200 ml of CSM (-Ura) + YNB medium supplemented with 2% inducer galactose. The culture was maintained at 30°C for 18 hours (until an O.D. between 1.0 and 2.0). The yeasts were centrifuged at 3,000 radiant power measurement, 5 minutes at 4°C, were washed once with distilled water, and the pellet was resuspended in lysis buffer (TEN: Tris 10 mM, pH 8.0; NaCl 150 mM; EDTA 1 mM) + 2X protease inhibitors (Compl Roche). A volume of glass beads having a size of about 425-600 microns (Sigma) were added for the lysis. This mixture was subjected to vigorous vortex stirring for 30 seconds 4 times, with 30-second intervals, and at 4°C. After this, the soluble fraction was recovered by centrifuging the lysis mixture at 13,000 rpm for 15 minutes at 4°C. This sample was subjected to fractioning on a sucrose gradient according to a previously described protocol (Lombardo, E., Maraver, A., Castón, J. R., Rivera, J., Fernández-Arias, A., Serrano, A., Carrascosa, J. L. & Rodríguez, J. F. (1999). VP1, the putative RNA-

dependent RNA polymerase of infectious bursal disease virus, forms complexes with the capsid protein VP3, leading to efficient encapsidation into virus-like particles. *Journal of Virology* 73, 6973-6983). The samples obtained after fractioning as well as a sample of the starting material were analyzed by means of sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) [Current Protocols in Molecular Biology] and immunodetection by Western blot (Figure 4A) using anti-pVP2 and anti-VP3 sera [Current Protocols in Molecular Biology]. As is shown in Figure 4A, the Western blot showed the presence of bands, with the predicted molecular mass corresponding to the pVP2 (48 kDa) and VP3-GFP (61 kDa) proteins, as well as other immunoreactive bands of a smaller size probably produced by proteolytic degradation both in the initial sample and in the different fractions of the gradient. These results reliably showed the correct expression of both polypeptides in the *S. cerevisiae* culture transformed with the plasmid pESCURA/pVP2-VP3. Then, the different fractions of the gradient were analyzed by means of TEM as has been previously described (Lombardo, E., Maraver, A., Castón, J. R., Rivera, J., Fernández-Arias, A., Serrano, A., Carrascosa, J. L. & Rodríguez, J. F. (1999). VP1, the putative RNA-dependent RNA polymerase of infectious bursal disease virus, forms complexes with the capsid protein VP3, leading to efficient encapsidation into virus-like particles. *Journal of Virology* 73, 6973-6983). As is shown in Figure 4B, the TEM analysis of the fractions of the gradient showed the existence of IBDV VLPs in the top fractions of the gradient. These VLPs, VLPs(-VP4), have a diameter of 65-70 nm and a polygonal contour that is indistinguishable from the IBDV VLPs obtained in other expression systems (Figure 4C).

CLAIMS

1. An empty capsid of the infectious bursal disease virus (IBDV), VLP(-VP4), characterized in that it is constituted by assembly of only IBDV pVP2 proteins and IBDV
5 VP3 proteins.
2. A nucleic acid characterized in that its nucleotide sequence is constituted by (i) a nucleotide sequence comprising the open reading frame corresponding to the IBDV pVP2 protein and (ii) a nucleotide sequence comprising the open reading frame corresponding to
10 the IBDV VP3 protein.
3. A gene construct comprising a nucleic acid according to claim 2.
4. An expression system selected from:
- 15 a) an expression system comprising (i) a gene construct comprising the open reading frame corresponding to the IBDV pVP2 protein, operatively bound to transcription, and optionally translation, control elements, and (ii) a gene construct comprising the open reading frame corresponding to
20 the IBDV VP3 protein, operatively bound to transcription, and optionally translation, control elements; and
- b) an expression system comprising a gene construct according to claim 3, operatively bound to transcription, and optionally translation, control
25 elements.
5. An expression system according to claim 4, said expression system being selected from plasmids, bacmids, yeast artificial chromosomes (YACs), bacteria artificial chromosomes (BACs), bacteriophage P1-based artificial chromosomes (PACs), cosmids,
30 and viruses, which, optionally, contain a heterologous replication origin.
6. A host cell containing a nucleic acid according to claim 2, or a gene construct according to claim 3, or an expression system according to anyone of claims 4 or 5.

7. A host cell that is transformed, transfected or infected with an expression system according to anyone of claims 4 or 5.

5 8. Host cell according to anyone of claims 6 or 7, said cell being an insect cell or a yeast.

9. A process for the production of empty capsids of the infectious bursal disease virus (IBDV), VLPs(-VP4), according to claim 1, comprising culturing a host cell according to
10 anyone of claims 6 to 8, and if so desired, recovering said empty IBDV capsids.

10. Process according to claim 9, wherein said host cell is an insect cell, comprising the steps of:

15 a) preparing an expression system selected from:

- an expression system constituted by a recombinant baculovirus containing a gene construct according to claim 3, operatively bound to transcription, and optionally translation, control elements; and

20

- an expression system constituted by (i) a recombinant baculovirus containing a gene construct comprising the open reading frame corresponding to the IBDV pVP2 protein, and (ii) a recombinant baculovirus containing a gene construct comprising the open reading
25 frame corresponding to the IBDV VP3 protein;

25

b) infecting insect cells with said expression system prepared in step a);

c) culturing the infected insect cells obtained in step b) under conditions
30 allowing the expression of recombinant proteins and their assembly for forming empty IBDV capsids, VLPs(-VP4); and

- d) if so desired, isolating and optionally purifying said IBDV empty capsids, VLPs(-VP4).

11. Process according to claim 9, wherein said host cell is a yeast, comprising the
5 steps of:

- a) preparing an expression system constituted by a plasmid containing a gene construct according to claim 3;
- 10 b) transforming yeast cells with said expression system prepared in step a);
- c) culturing the transformed yeasts obtained in step b) under conditions allowing the expression of recombinant proteins and their assembly to form empty IBDV capsids, VLPs(-VP4); and
- 15 d) if so desired, isolating and optionally purifying the empty IBDV capsids, VLPs(-VP4).

12. The use of a gene expression system according to anyone of claims 4 or 5 for
20 producing and obtaining empty IBDV capsids, VLPs(-VP4), according to claim 1.

13. The use of empty capsids of the infectious bursal disease virus (IBDV), VLPs(-VP4), according to claim 1 in the manufacture of a medicament.

25 14. Use according to claim 13, wherein said medicament is a vaccine against the avian disease called infectious bursal disease.

15. Use according to claim 13, wherein said medicament is a gene therapy vector.

30 16. A vaccine comprising a therapeutically effective amount of empty IBDV capsids, VLPs(-VP4), according to claim 1, optionally together with one or more pharmaceutically acceptable adjuvants and/or vehicles.

17. Vaccine according to claim 16 to protect birds from the infection caused by the infectious bursal disease virus (IBDV).

18. Vaccine according to claim 17, wherein said birds are selected from the group
5 formed by chickens, turkeys, geese, ganders, pheasants, quails and ostriches.

19. Vaccine to protect chickens from the infection caused by the infectious bursal disease virus (IBDV) comprising a therapeutically effective amount of empty IBDV capsids, VLPs(-VP4), according to claim 1, optionally together with one or more pharmaceutically
10 acceptable adjuvants and/or vehicles.




Fig. 1




Fig. 2




Fig. 3




Fig. 4

SEQUENCE LISTING

<110> CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS
<110> BIONOSTRA, S.L.

<120> EMPTY CAPSIDS (VLPs(-VP4)) OF THE INFECTIOUS BURSAL DISEASE VIRUS
(IBDV), OBTAINMENT PROCESS AND APPLICATIONS

<130> P1392PC

<150> ES P200400121
<151> 2004-01-21 (January 21, 2004)

<160> 10
<170> PatentIn version 3.1

<210> 1
<211> 35
<212> DNA
<213> Artificial sequence

<220> Synthetic DNA
<223> Oligo I primer

<400> 1
gcgcagatct atgacaaacc tgcagatca aaccc 35

<210> 2
<211> 34
<212> DNA
<213> Artificial sequence

<220> Synthetic DNA
<223> Oligo II primer

<400> 2
gcgcaagctt aggcgagagt cagctgcctt atgc 34

<210> 3
<211> 7595
<212> DNA
<213> Artificial sequence

<220>
<223> Plasmid pFBD/pVP2-his-VP3

<221> promoter
<222> (157)..(285)
<223> Promotor ppolh

<221> CDS
<222> (291)..(1289)
<223> pVP2 ORF

agg ctt ggt gac ccc att ccc gca ata ggg ctt gac cca aaa atg gta	872
Arg Leu Gly Asp Pro Ile Pro Ala Ile Gly Leu Asp Pro Lys Met Val	
180 185 190	
gcc aca tgt gac agc agt gac agg ccc aga gtc tac acc ata act gca	920
Ala Thr Cys Asp Ser Ser Asp Arg Pro Arg Val Tyr Thr Ile Thr Ala	
195 200 205 210	
gcc gat gat tac caa ttc tca tca cag tac caa cca ggt ggg gta aca	968
Ala Asp Asp Tyr Gln Phe Ser Ser Gln Tyr Gln Pro Gly Gly Val Thr	
215 220 225	
atc aca ctg ttc tca gcc aac att gat gcc atc aca agc ctc agc gtt	1016
Ile Thr Leu Phe Ser Ala Asn Ile Asp Ala Ile Thr Ser Leu Ser Val	
230 235 240	
ggg gga gag ctc gtg ttt cga aca agc gtc cac ggc ctt gta ctg ggc	1064
Gly Gly Glu Leu Val Phe Arg Thr Ser Val His Gly Leu Val Leu Gly	
245 250 255	
gcc acc atc tac ctc ata ggc ttt gat ggg aca acg gta atc acc agg	1112
Ala Thr Ile Tyr Leu Ile Gly Phe Asp Gly Thr Thr Val Ile Thr Arg	
260 265 270	
gct gtg gcc gca aac aat ggg ctg acg acc ggc acc gac aac ctt atg	1160
Ala Val Ala Ala Asn Asn Gly Leu Thr Thr Gly Thr Asp Asn Leu Met	
275 280 285 290	
cca ttc aat ctt gtg att cca aca aac gag ata acc cag cca atc aca	1208
Pro Phe Asn Leu Val Ile Pro Thr Asn Glu Ile Thr Gln Pro Ile Thr	
295 300 305	
tcc atc aaa ctg gag ata gtg acc tcc aaa agt ggt ggt cag gca ggg	1256
Ser Ile Lys Leu Glu Ile Val Thr Ser Lys Ser Gly Gly Gln Ala Gly	
310 315 320	
gat cag atg tca tgg tcg gca aga ggg agc cta gcagtgcgca tccatggtgg	1309
Asp Gln Met Ser Trp Ser Ala Arg Gly Ser Leu	
325 330	
caactatcca ggggccctcc gtcccgtcac gctagtggcc tacgaaagag tggcaacagg	1369
atccgtcggt acggtcgctg ggggtgagcaa ctccgagctg atcccaaata ctgaactagc	1429
aaagaacctg gttacagaat acggccgatt tgaccagga gccatgaact acacaaaatt	1489
gatactgagt gagagggacc gtcttgat caagaccgtc tggccaacaa gggagtacac	1549
tgactttcgt gaatacttca tggaggtggc cgacctcaac tctcccctga agattgcagg	1609
agcattcggc ttcaaagaca taatccgggc cataaggagg atagctgtgc cgggtgtctc	1669
cacattgttc ccacctgccg ctcccctagc ccatgcaatt ggggaagggtg tagactacct	1729
gctgggcat gagggccagg ccgcttcagg aactgctcga gccgcgtcag gaaaagcaag	1789
agctgcctca ggccgcataa ggcagctgac tctcgcctaa gcttgctcag aagtactaga	1849
ggatcataat cagccatacc acattttagt aggttttact tgctttaaaa aacctccac	1909

acctccccct gaacctgaaa cataaaatga atgcaattgt tgttgtaac ttgtttattg 1969
cagcttataa tggttacaaa taaagcaata gcatcacaaa ttccacaaat aaagcatttt 2029
tttcaactgca ttctagttgt ggtttgtcca aactcatcaa tgtatcttat catgtctgga 2089
tctgatcact gcttgagcct aggagatccg aaccagataa gtgaaatcta gttccaaact 2149
attttgtcat ttttaatttt cgtattagct tacgacgcta caccagttc ccatctattt 2209
tgtcactctt ccctaaataa tccttaaaaa ctccatttcc acccctccca gttcccaact 2269
attttgtccg cccacagcgg ggcatttttc ttctgttat gtttttaatc aaacatcctg 2329
ccaactccat gtgacaaacc gtcatcttcg gctacttttt ctctgtcaca gaatgaaaat 2389
ttttctgtca tctcttcggt attaatgttt gtaattgact gaatatcaac gcttatttgc 2449
agcctgaatg gcgaatggga cgcgcctgt agcggcgcat taagcgcggc ggggtgtggtg 2509
gttacgcgca gcgtgaccgc tacacttgcc agcgcctag cgcccgctcc ttctgcttcc 2569
ttcccttcc tctctgccac gttcgccggc ttcccccgtc aagctctaaa tcgggggctc 2629
cctttagggt tccgatttag tgctttacgg cacctcgacc ccaaaaaact tgattagggt 2689
gatggttcac gtagtgggcc atcgccctga tagacggttt ttcgcccttt gacgttgag 2749
tccacgttct ttaatagtgg actcttggtc caaactggaa caaactcaa ccctatctcg 2809
gtctattctt ttgatttata agggattttg cagatttcgg cctattggtt aaaaaatgag 2869
ctgatttaac aaaaatttaa cgcgaatttt aacaaaatat taacgtttac aatttcaggt 2929
ggcacttttc ggggaaatgt gcgcggaacc cctatttggt tatttttcta aatacattca 2989
aatatgtatc cgctcatgag acaataaacc tgataaatgc ttcaataata ttgaaaaagg 3049
aagagtatga gtattcaaca ttccogtgc gcccttattc ctttttttgc ggcattttgc 3109
cttctgttt ttgtcacc agaaacgtg gtgaaagtaa aagatgctga agatcagttg 3169
ggtgacagag tgggttacat cgaactggat ctcaacagcg gtaagatcct tgagagtttt 3229
cgccccgaag aacgttttcc aatgatgagc acttttaaag ttctgctatg tggcgcggt 3289
ttatcccgta ttgacgccg gcaagagcaa ctcggtcgcc gcatacacta ttctcagaat 3349
gacttggtt agtactcacc agtcacagaa aagcatctta cggatggcat gacagtaaga 3409
gaattatgca gtgtgccat aaccatgagt gataaactg cgccaactt acttctgaca 3469
acgatcgag gaccgaagga gctaaccgct tttttgcaca acatggggga tcatgtaact 3529
cgccctgac gttgggaacc ggagctgaat gaagccatac caaacgacga gcgtgacacc 3589
acgatgcctg tagcaatggc aacaocgtt cgcaaactat taactggcga actacttact 3649
ctagcttccc ggcaacaatt aatagactgg atggaggcgg ataaagttgc aggaccactt 3709

ctgcgctcgg cccttccggc tggctggttt attgctgata aatctggagc cggtgagcgt 3769
gggtctcgcg gtatcattgc agcactgggg ccagatggta agccctcccg tatcgtagtt 3829
atctacacga cggggagtca ggcaactatg gatgaacgaa atagacagat cgctgagata 3889
gggtgcctcac tgattaagca ttggtaactg tcagaccaag tttactcata tatacttttag 3949
attgatttaa aacttcattt ttaatttaaa aggatctagg tgaagatcct ttttgataat 4009
ctcatgacca aaatccctta acgtgagttt tcgttccact gagcgtcaga ccccgtagaa 4069
aagatcaaag gatcttcttg agatcctttt tttctgcgcg taatctgctg cttgcaaaca 4129
aaaaaaccac cgctaccagc ggtgggtttgt ttgccggatc aagagctacc aactcttttt 4189
ccgaaggtaa ctggcttcag cagagcgagc ataccaaata ctgtccttct agtgtagccg 4249
tagttaggcc accacttcaa gaactctgta gcaccgccta catacctcgc tctgctaate 4309
ctgttaccag tggctgctgc cagtggcgat aagtcgtgtc ttaccgggtt ggactcaaga 4369
cgatagttac cggataaggc gcagcggtcg ggctgaacgg ggggttcgtg cacacagccc 4429
agcttggagc gaacgaccta caccgaactg agatacctac agcgtgagca ttgagaaagc 4489
gccacgcttc ccgaaggag aaaggcggac aggtatccgg taagcggcag ggtcggaaaca 4549
ggagagcgca cgaggagct tccaggggga aacgcctggg atctttatag tcctgtcggg 4609
tttcgccacc tctgacttga gcgtcgattt ttgtgatgct cgtcaggggg gcggagccta 4669
tggaaaaaacg ccagcaacgc ggccctttta cggttcctgg ccttttgctg gccttttct 4729
cacatgttct ttcctgcgtt atcccctgat tctgtggata accgtattac cgcctttgag 4789
tgagctgata ccgctcgccg cagccgaacg accgagcgca gcgagtcagt gagcgaggaa 4849
gcggaagagc gcctgatgcg gtattttctc cttacgcate tgtcgggtat ttcacaccgc 4909
agaccagccg cgtaacctgg caaaatcggg tacggttgag taataaatgg atgccctgcg 4969
taagcgggtg tgggaggaca ataaagtctt aaactgaaca aaatagatct aaactatgac 5029
aataaagtct taaactagac agaatagttg taaactgaaa tcagtccagt tatgctgtga 5089
aaaagcatac tggacttttg ttatggctaa agcaaactct tcattttctg aagtgcaaat 5149
tgcccgtcgt attaaagagg ggcgtggcca agggcatggg aaagactata ttcgcggcgt 5209
tgtgacaatt taccgaacaa ctccgcgcc gggaagccga tctcggcttg aacgaattgt 5269
taggtggcgg tacttgggtc gatatcaaag tgcactactt cttcccgtat gcccaacttt 5329
gtatagagag cactgcggg atcgtcaccg taatctgctt gcacgtagat cacataagca 5389
ccaagcgcgt tggcctcatg cttgaggaga ttgatgagcg cggtggaat gccctgcctc 5449
cgggtgctcgc cggagactgc gagatcatag atatagatct cactacgcgg ctgctcaaac 5509

ctgggcagaa cgtaagccgc gagagcgcca acaaccgctt cttggtcgaa ggcagcaagc 5569
gcgatgaatg tottactacg gagcaagtcc cggaggtaat cggagtccgg ctgatgttgg 5629
gagtaggtgg ctacgtctcc gaactcacga ccgaaaagat caagagcagc ccgcatggat 5689
ttgacttggt cagggccgag cctacatgtg cgaatgatgc ccatacttga gccacctaac 5749
tttgtttttag ggcgactgcc ctgctgcgta acatcgttgc tgctgcgtaa catcgttgct 5809
gtcccataac atcaaacatc gacccacggc gtaacgcgct tgctgcttgg atgcccgagg 5869
catagactgt acaaaaaaac agtcataaca agccatgaaa accgccactg cgccgttacc 5929
accgctgcgt tcggtcaagg ttctggacca gttgcgtgag cgcatacgt acttgatta 5989
cagtttacga accgaacagg cttatgtcaa ctgggttcgt gccttcatcc gttccacgg 6049
tgtgcgtcac ccggcaacct tgggcagcag cgaagtcgag gcatttctgt cctggctggc 6109
gaacgagcgc aaggtttcgg tctccacgca tcgtcaggca ttggcggcct tgctgttctt 6169
ctacggcaag gtgctgtgca cggatctgcc ctggcttcag gagatcggtg gacctcgcc 6229
gtcgcggcgc ttgccggtgg tgctgacccc ggatgaagtg gttcgcaccc tcggttttct 6289
ggaaggcgag catcgtttgt tcgcccagga ctctagctat agttctagtg gttggcctac 6349
gtaccgtag tggctatggc agggcttgcc gccccgacgt tggtgcgag ccctgggcct 6409
tcacccgaac ttgggggttg ggggtgggaa aaggaaagaaa cgcgggcgta ttggtccaa 6469
tggggtctcg gtggggtatc gacagagtgc cagccctggg accgaacccc gcgtttatga 6529
acaaacgacc caacaccggt gcgttttatt ctgtcttttt attgccgtca tagcgcggt 6589
tccttccggt attgtctcct tccgtgttcc agttagcctc ccccatctcc cggtaccgca 6649
tgccctgaga ctgcaggctc tagattcgaa agcggccgag actagtgagc tcgtcgacgt 6709
aggcctttga attccggatc ctactcaag gtcctcatca gagacggtcc tgatccagcg 6769
gccagccga ccagggggtc tctgtgttgg agcattgggt tttggcttgg gctttggtag 6829
agcccgctg ggattgcgat gcttcatctc catcgcagtc aagagcagat ctttcatctg 6889
ttcttggttt gggccacgtc catggttgat tcatagact ttggcaactt cgtctatgaa 6949
agcttggggt ggctctgcct gtcctggagc cccgtagatc gacgtagctg cccttaggat 7009
ttgttcttct gatgccaaac ggctcttctc tgcatgcacg tagtctagat agtctcgtt 7069
tgggtccggt atttctcggt tgttctgcca gtactttacc tggcctgggc ttggccctcg 7129
gtgccattg agtgctaccc attctggtgt tgcaaagtag atgcccatgg tctccatctt 7189
ctttgagatc cgtgtgtctt tttccctctg tgcttctctt ggtgtggggc ccgagcctc 7249
cactccgtag cctgctgtcc cgtacttggc cctttgcgac ttgctgcctg cttgtggtgc 7309

gtttgcaaga aaatttcgca tccgatgggc gttcgggtcg ctgagtgcga agttggccat 7369
 gtcagtcaca atccattct cttccagcca catgaacaca ctgagtgcag attggaatag 7429
 tgggtccacg ttggctgctg cttccattgc tctgacggca ctctcgagtt cgggggtctc 7489
 tttgaactct gatgcagcca tggcgccctg aaaatacagg ttttcggtcg ttgggatatc 7549
 gtaatcgtga tgggtgatggt gatggtagta cgacatgggt tcggac 7595

<210> 4
 <211> 333
 <212> PRT
 <213> Artificial sequence

<220>
 <223> pVP2-his-VP3 protein

<400> 4
 Met Thr Asn Leu Ser Asp Gln Thr Gln Gln Ile Val Pro Phe Ile Arg
 1 5 10 15
 Ser Leu Leu Met Pro Thr Thr Gly Pro Ala Ser Ile Pro Asp Asp Thr
 20 25 30
 Leu Glu Lys His Thr Leu Arg Ser Glu Thr Ser Thr Tyr Asn Leu Thr
 35 40 45
 Val Gly Asp Thr Gly Ser Gly Leu Ile Val Phe Phe Pro Gly Phe Pro
 50 55 60
 Gly Ser Ile Val Gly Ala His Tyr Thr Leu Gln Gly Asn Gly Asn Tyr
 65 70 75 80
 Lys Phe Asp Gln Met Leu Leu Thr Ala Gln Asn Leu Pro Ala Ser Tyr
 85 90 95
 Asn Tyr Cys Arg Leu Val Ser Arg Ser Leu Thr Val Arg Ser Ser Thr
 100 105 110
 Leu Pro Gly Gly Val Tyr Ala Leu Asn Gly Thr Ile Asn Ala Val Thr
 115 120 125
 Phe Gln Gly Ser Leu Ser Glu Leu Thr Asp Val Ser Tyr Asn Gly Leu
 130 135 140
 Met Ser Ala Thr Ala Asn Ile Asn Asp Lys Ile Gly Asn Val Leu Val
 145 150 155 160
 Gly Glu Gly Val Thr Val Leu Ser Leu Pro Thr Ser Tyr Asp Leu Gly
 165 170 175
 Tyr Val Arg Leu Gly Asp Pro Ile Pro Ala Ile Gly Leu Asp Pro Lys
 180 185 190
 Met Val Ala Thr Cys Asp Ser Ser Asp Arg Pro Arg Val Tyr Thr Ile
 195 200 205

Thr Ala Ala Asp Asp Tyr Gln Phe Ser Ser Gln Tyr Gln Pro Gly Gly
 210 215 220
 Val Thr Ile Thr Leu Phe Ser Ala Asn Ile Asp Ala Ile Thr Ser Leu
 225 230 235 240
 Ser Val Gly Gly Glu Leu Val Phe Arg Thr Ser Val His Gly Leu Val
 245 250 255
 Leu Gly Ala Thr Ile Tyr Leu Ile Gly Phe Asp Gly Thr Thr Val Ile
 260 265 270
 Thr Arg Ala Val Ala Ala Asn Asn Gly Leu Thr Thr Gly Thr Asp Asn
 275 280 285
 Leu Met Pro Phe Asn Leu Val Ile Pro Thr Asn Glu Ile Thr Gln Pro
 290 295 300
 Ile Thr Ser Ile Lys Leu Glu Ile Val Thr Ser Lys Ser Gly Gly Gln
 305 310 315 320
 Ala Gly Asp Gln Met Ser Trp Ser Ala Arg Gly Ser Leu
 325 330

<210> 5
 <211> 35
 <212> DNA
 <213> Artificial sequence

<220> Synthetic DNA
 <223> Oligo III primer

<400> 5
 gcgcagatct atgacaaacc tgtcagatca aaccc

35

<210> 6
 <211> 34
 <212> DNA
 <213> Artificial sequence

<220> Synthetic DNA
 <223> Oligo IV primer

<400> 6
 gcgcaagctt aggcgagagt cagctgcctt atgc

34

<210> 7
 <211> 33
 <212> DNA
 <213> Artificial sequence

<220> Synthetic DNA
 <223> Oligo V primer

<400> 7
 gcgcgaattc gatggcatca gagttcaaag aga

33

<210> 8
 <211> 32
 <212> DNA
 <213> Artificial sequence

<220> Synthetic DNA
 <223> Oligo VI primer

<400> 8
 cgcgcatccc tcaaggctct catcagagac gg

32

<210> 9
 <211> 9600
 <212> DNA
 <213> Artificial sequence

<223> Plasmid pESCURA/pVP2-VP3-GFP

<221> promoter
 <222> (5649)..(5859)
 <223> Promoter GAL 1 (pVP2)

<221> promoter
 <222> (7402)..(8080)
 <223> Promoter GAL 2 (VP3-GFP)

<221> CDS
 <222> (8086)..(9597)
 <223> VP3-GFP ORF

<400> 9
 ggccgcacta gtatcgatgg attacaagga tgacgacgat aagatctgag ctcttaatta 60
 acaattcttc gccagagggt tggtaagtc tccaatcaag gttgtcggct tgtctacctt 120
 gccagaaatt tacgaaaaga tggaaaaggg tcaaatcggt ggtagatacg ttgttgacac 180
 ttctaaataa gcgaatttct tatgatttat gatttttatt attaaataag ttataaaaaa 240
 aataagtgtg tacaaaattt aaagtgactc ttaggtttta aaacgaaaat tcttattctt 300
 gagtaactct ttctgttagg tcaggttget ttctcaggta tagcatgagg tcgctccaat 360
 tcagctgcat taatgaatcg gccaacgcgc ggggagaggc ggtttgcgta ttgggcgctc 420
 ttccgcttcc tcgctcactg actcgtcgcg ctcggtcggt cggctgcggc gagcgggtatc 480
 agctcactca aaggcggtta tacggttatc cacagaatca ggggataacg caggaaagaa 540
 catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tgctggcggt 600
 ttccataggt ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa gtcagagggtg 660
 gcgaaacccg acaggactat aaagatacca ggcgtttccc cctggaagct ccctcgtgcg 720
 ctctcctggt ccgaccctgc cgcttacggg atacctgtcc gcctttctcc cttcggaag 780
 cgtggcgctt tctcatagct cacgctgtag gtatctcagt tcggtgtagg tcgttcgctc 840

caagctgggc tgtgtgcacg aaccccccg tccagccgac cgctgcgcct tatccggtaa 900
ctatcgtctt gagtccaacc cggtaagaca cgacttatcg ccaactggcag cagccactgg 960
taacaggatt agcagagcga ggtatgtagg cgggtgtaca gagttcttga agtgggtggc 1020
taactacggc tacactagaa ggacagtatt tggatatctgc gctctgctga agccagttac 1080
cttcggaaaa agagttggta gctcttgatc cggcaaacia accaccgctg gtagcgggtg 1140
tttttttgtt tgcaagcagc agattacgcg cagaaaaaaa ggatctcaag aagatccttt 1200
gactttttct acgggggtctg acgctcagtg gaacgaaaac tcacgttaag ggatttttgt 1260
catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaaaat gaagttttaa 1320
atcaatctaa agtatatatg agtaaaacttg gtctgacagt taccaatgct taatcagtga 1380
ggcacctatc tcagcgatct gtctatctcg ttcattcata gttgcctgac tcccgcgtgt 1440
gtagataact acgatacggg agggcttacc atctggcccc agtgctgcaa tgataccgcg 1500
agaccacgc tcaccggctc cagatttatc agcaataaac cagccagccg gaagggccga 1560
gcgcagaagt ggtcctgcaa ctttatccgc ctccatccag tctattaatt gttgccggga 1620
agctagagta agtagttcgc cagttaatag tttgcgcaac gttgttgcca ttgctacagg 1680
catcgtggtg tcacgctcgt cgtttggtat ggcttcattc agctccggtt cccaacgatc 1740
aaggcgagtt acatgatccc ccatgttgtg caaaaaagcg gttagctcct tcggtcctcc 1800
gatcgttgtc agaagtaagt tggccgcagt gttatcactc atggttatgg cagcactgca 1860
taattctctt actgtcatgc catccgtaag atgcttttct gtgactgggt agtactcaac 1920
caagtcattc tgagaatagt gtatgcggcg accgagttgc tcttgcccg cgtaataacg 1980
ggataatacc gcgccacata gcagaacttt aaaagtgtc atcattggaa aacgttcttc 2040
ggggcgaaaa ctctcaagga tcttaccgct gttgagatcc agttcgatgt aaccactcg 2100
tgacccaac tgatcttcag catcttttac tttcaccagc gtttctgggt gagcaaaaac 2160
aggaaggcaa aatgccgcaa aaaagggaat aagggcgaca cggaaatgtt gaatactcat 2220
actcttctt tttcaatatt attgaagcat ttatcagggt tattgtctca tgagcggata 2280
catatttgaa tgtatttaga aaaataaaca aataggggtt ccgcgacat tccccgaaa 2340
agtgcacact gaacgaagca tctgtgcttc atttttaga acaaaaatgc aacgcgagag 2400
cgctaatttt tcaacaaaag aatctgagct gcatttttac agaacagaaa tgcaacgcga 2460
aagcgtatatt ttaccaacga agaactctgt cttcattttt gtaaaacaaa aatgcaacgc 2520
gagagcgcta atttttcaaa caaagaatct gagctgcatt ttacagaaac agaaatgcaa 2580
cgcgagagcg ctattttacc aacaaagaat ctatacttct tttttgttct acaaaaatgc 2640

atccccgagag cgctatTTTT ctaacaaagc atcttagatt actTTTTTtc tcctttgtgc 2700
gctctataat gcagtcctct gataactttt tgcactgtag gtccgttaag gttagaagaa 2760
ggctactttg gtgtctatTT tctcttccat aaaaaagcc tgactccact tcccgcgttt 2820
actgattact agcgaagctg cgggtgcatt ttttcaagat aaaggcatcc ccgattatat 2880
tctataccga tgtggattgc gcatactttg tgaacagaaa gtgatagcgt tgatgattct 2940
tcattgggtca gaaaattatg aacggtttct tctatTTTgt ctctatatac tacgtatagg 3000
aaatgtttac attttcgtat tgttttcgat tcaactctatg aatagttctt actacaattt 3060
ttttgtctaa agagtaatac tagagataaa cataaaaaat gtagaggtcg agtttagatg 3120
caagttcaag gagcgaaagg tggatgggta gggtatatag ggatatagca cagagatata 3180
tagcaaagag atacttttga gcaatgtttg tggaagcggg attcgcaata ttttagtagc 3240
tcgttacagt ccggtgcgtt tttggTTTT tgaaagtgcg tcttcagagc gcttttggtt 3300
ttcaaaagcg ctctgaagtt cctatacttt ctagagaata ggaacttcgg aataggaact 3360
tcaaagcgtt tccgaaaacg agcgcttccg aaaatgcaac gcgagctgcg cacatacagc 3420
tcaactgttca cgtcgcacct atatctgcgt gttgcctgta tatatatata catgagaaga 3480
acggcatagt gcgtgtttat gcttaaatgc gtacttatat gcgtctattt atgtaggatg 3540
aaaggtagtc tagtacctcc tgtgatatta tccattcca tgcggggtat cgtatgcttc 3600
cttcagcaact acccttttagc tgttctatat gctgccactc ctcaattgga ttagtctcat 3660
ccttcaatgc tatcatttcc tttgatattg gatcatacta agaaaccatt attatcatga 3720
cattaacctt taaaaatagg cgtatcacga ggccctttcg tctcgcgcgt ttcggtgatg 3780
acggtgaaaa cctctgacac atgcagctcc cggagacggg cacagcttgt ctgtaagcgg 3840
atgccgggag cagacaagcc cgtcagggcg cgtcagcggg tgttggcggg tgcggggcgt 3900
ggcttaacta tgcggcatca gagcagattg tactgagagt gcaccatacc acagcttttc 3960
aattcaatlc atcattTTTT ttttattctt ttttttgatt tcggtttctt tgaaatTTTT 4020
ttgattcggg aatctccgaa cagaaggaag aacgaaggaa ggagcacaga cttagattgg 4080
tatatatacg catatgtagt gttgaagaaa catgaaattg ccaggtattc ttaacccaac 4140
tgcacagAAC aaaaacctgc aggaaacgaa gataaatcat gtcgaaagct acatataagg 4200
aacgtgctgc tactcatcct agtcctgttg ctgccaagct atttaatatc atgcacgaaa 4260
agcaacaaa cttgtgtgct tcattggatg ttcgtaccac caaggaatta ctggagttag 4320
ttgaagcatt aggtcccaaa atttgTTTtac taaaaacaca tgtggatatc ttgactgatt 4380
ttccatgga gggcacagtt aagccgctaa aggcattatc cgccaagtac aattTTTTtac 4440

tcttogaaga cagaaaaattt gctgacattg gtaatacagt caaattgcag tactctgcgg 4500
gtgtatacag aatagcagaa tgggcagaca ttacgaatgc acacgggtgtg gtgggccag 4560
gtattgttag cggtttgaag caggcggcag aagaagtaac aaaggaacct agaggccttt 4620
tgatgttagc agaattgtca tgcaagggt .ccctatctac tggagaatat actaagggt 4680
ctgttgacat tgcaagagc gacaaagatt ttgttatcgg ctttattgct caaagagaca 4740
tgggtggaag agatgaagggt tacgattggt tgattatgac acccgggtgtg gggttagatg 4800
acaagggaga cgcattgggt caacagtata gaaccgtgga tgatgtggtc tctacaggat 4860
ctgacattat tattgttga agaggactat ttgcaaagg aagggatgct aaggtagagg 4920
gtgaacgtta cagaaaagca ggctgggaag catatttgag aagatgcggc cagcaaaact 4980
aaaaaactgt attataagta aatgcatgta tactaaactc acaaattaga gcttcaattt 5040
aattatatca gttattacc tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa 5100
taccgcatca ggaaattgta aacgttaata tttgttaaa attcgcgtta aatttttgtt 5160
aatcagctc attttttaac caataggccg aaatcgcaa aatcccttat aaatcaaaag 5220
aatagaccga gatagggttg agtgtgttc cagtttgaa caagagtcca ctattaaaga 5280
acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc ccactacgtg 5340
aaccatcacc ctaatcaagt ttttgggggt cgagggtccg taaagcacta aatcggaacc 5400
ctaaaggag ccccgattt agagcttgac ggggaaagcc gggaacgtg gcgagaaagg 5460
aagggaagaa agcgaaagga gcgggcgcta gggcgctggc aagtgtagcg gtcacgctgc 5520
gcgtaaccac cacaccgcc gcgcttaatg cgccgtaca gggcgcgctc cgccattcgc 5580
cattcaggct gcgcaactgt tgggaagggc gatcggtgag ggcctcttcg ctattacgcc 5640
agctggatct tcgagcgtcc caaaaacctc tcaagcaagg ttttcagtat aatgttacat 5700
gcgtacacgc gtctgtacag aaaaaaaga aaaatttgaa atataaataa cgttcttaat 5760
actaacataa ctataaaaaa ataaataggg acctagactt caggttgtct aactccttcc 5820
ttttcggtta gagcgatct tagctagccg cggtagcaag ctaggcgag agtcagctgc 5880
cttatgcggc ctgaggcagc tottgctttt cctgacgcgg ctcgagcagt tcctgaagcg 5940
gcctgggcct catcgccag caggtagtct acaccttccc caattgcatg ggctagggga 6000
ggggcagggtg ggaacaatgt ggagaccacc ggcacagcta tcctccttat ggcccggtt 6060
atgtctttga agccgaatgc tcctgcaatc ttcaggggag agttgaggtc ggccacctcc 6120
atgaagtatt cacgaaagtc agtgactcc cttgttgcc agacggtctt gatgccaaga 6180
cggtccctct cactcagtat caattttgtg tagttcatgg ctctgggtc aaatcggccg 6240

tattctgtaa ccaggttctt tgctagttca ggatttggga tcagctcgaa gttgctcacc 6300
ccagcgaccg taacgacgga tcctgttgcc actctttcgt aggccactag cgtgacggga 6360
cggagggccc ctggatagtt gccaccatgg atcgtcactg ctaggctccc tcttgccgac 6420
catgacatct gatccctgc ctgaccacca cttttggagg tcactatctc cagtttgatg 6480
gatgtgattg gctgggttat ctcgtttggt ggaatcacia gattgaatgg cataaggttg 6540
tcggtgccgg tcgtcagccc attgtttgcg gccacagccc tgggtgattac cgttgtecca 6600
tcaaagccta tgaggtagat ggtggcgccc agtacaaggc cgtggacgct tgttcgaaac 6660
acgagctctc ccccaacgct gaggcttggt atggcatcaa tgttggtga gaacagtgtg 6720
attgttacct cacctggttg gtactgtgat gagaattggt aatcatcggc tgcagttatg 6780
gtgtagactc tgggcctgct actgctgtca catgtggcta ccatttttgg gtcaagccct 6840
attgcgggaa tggggtcacc aagcctcaca taccgaagat catatgatgt gggtaagctg 6900
aggacggtga ccccttcccc tactaggacg ttcccaattt tgcgttgat gttggctggt 6960
gcagacatca acccattgta gctaacatct gtcagttcac tcaggcttcc ttggaaggctc 7020
acggcgttta tgggtccggt tagtgcataa acgccaccag gaagtgtgct tgacctcact 7080
gtgagactcc gactcactag cctgcagtag ttgtaactgg ccggtagggt ctgggcagtc 7140
aggagcatct gatcgaaact gtagttccca ttgccctgca gtgtgtagtg agcaccacaca 7200
attgagccag ggaatccagg gaaaaagaca attagccctg accctgtgtc cccacagtc 7260
aaattgtagg tcgaggtctc tgacctgaga gtgtgcttct ccagggtgtc gtccggaatg 7320
gacgccggtc cggttggttg catcagaagg ctccgtatga acggaacaat ctgctgggtt 7380
tgatctgaca ggtttgatcat agatccgggg tttttctctc ttgacgttaa agtatagagg 7440
tatattaaca attttttggt gatactttta ttacatttga ataagaagta atacaaaccg 7500
aaaatgttga aagtattagt taaagtgggt atgcagtttt tgcatttata tatctgttaa 7560
tagatcaaaa atcatcgctt cgctgattaa ttaccccaga aataaggcta aaaaactaat 7620
cgcattatca tcctatgggt gtaatttga ttcgttcatt tgaaggtttg tggggccagg 7680
ttactgcaa tttttcctct tcataacat aaaagctagt attgtagaat ctttattgtt 7740
cggagcagtg cggcgcgagg cacatctgcg tttcaggaaac gcgaccggtg aagacgagga 7800
cgcacggagg agagtcttcc ttcggagggc tgtcaccgcg tcggcggtt ctaatccgta 7860
cttcaatata gcaatgagca gttaagcgta ttactgaaag ttccaaagag aagggttttt 7920
taggctaaga taatggggct ctttacattt ccacaacata taagtaagat tagatatgga 7980
tatgtatatg gatatgtata tgggtggtaat gccatgtaat atgattatta aacttctttg 8040

cg	tcc	atc	cca	aaaaaaa	agt	aaga	attttt	gaaa	attcga	attcg	atg	gct	gca	tca	8097	
													Met	Ala	Ala	Ser
													1			
gag	ttc	aaa	gag	acc	ccc	gaa	ctc	gag	agt	gcc	gtc	aga	gca	atg	gaa	8145
Glu	Phe	Lys	Glu	Thr	Pro	Glu	Leu	Glu	Ser	Ala	Val	Arg	Ala	Met	Glu	
5					10					15					20	
gca	gca	gcc	aac	gtg	gac	cca	cta	ttc	caa	tct	gca	ctc	agt	gtg	ttc	8193
Ala	Ala	Ala	Asn	Val	Asp	Pro	Leu	Phe	Gln	Ser	Ala	Leu	Ser	Val	Phe	
				25					30					35		
atg	tgg	ctg	gaa	gag	aat	ggg	att	gtg	act	gac	atg	gcc	aac	ttc	gca	8241
Met	Trp	Leu	Glu	Glu	Asn	Gly	Ile	Val	Thr	Asp	Met	Ala	Asn	Phe	Ala	
			40					45					50			
ctc	agc	gac	ccg	aac	gcc	cat	cgg	atg	cga	aat	ttt	ctt	gca	aac	gca	8289
Leu	Ser	Asp	Pro	Asn	Ala	His	Arg	Met	Arg	Asn	Phe	Leu	Ala	Asn	Ala	
			55				60					65				
cca	caa	gca	ggc	agc	aag	tcg	caa	agg	gcc	aag	tac	ggg	aca	gca	ggc	8337
Pro	Gln	Ala	Gly	Ser	Lys	Ser	Gln	Arg	Ala	Lys	Tyr	Gly	Thr	Ala	Gly	
	70					75					80					
tac	gga	gtg	gag	gct	cgg	ggc	ccc	aca	cca	gag	gaa	gca	cag	agg	gaa	8385
Tyr	Gly	Val	Glu	Ala	Arg	Gly	Pro	Thr	Pro	Glu	Glu	Ala	Gln	Arg	Glu	
85					90					95					100	
aaa	gac	aca	cgg	atc	tca	aag	aag	atg	gag	acc	atg	ggc	atc	tac	ttt	8433
Lys	Asp	Thr	Arg	Ile	Ser	Lys	Lys	Met	Glu	Thr	Met	Gly	Ile	Tyr	Phe	
				105					110					115		
gca	aca	cca	gaa	tgg	gta	gca	ctc	aat	ggg	cac	cga	ggg	cca	agc	cca	8481
Ala	Thr	Pro	Glu	Trp	Val	Ala	Leu	Asn	Gly	His	Arg	Gly	Pro	Ser	Pro	
			120					125					130			
ggc	cag	gta	aag	tac	tgg	cag	aac	aaa	cga	gaa	ata	ccg	gac	cca	aac	8529
Gly	Gln	Val	Lys	Tyr	Trp	Gln	Asn	Lys	Arg	Glu	Ile	Pro	Asp	Pro	Asn	
		135					140					145				
gag	gac	tat	cta	gac	tac	gtg	cat	gca	gag	aag	agc	cgg	ttg	gca	tca	8577
Glu	Asp	Tyr	Leu	Asp	Tyr	Val	His	Ala	Glu	Lys	Ser	Arg	Leu	Ala	Ser	
	150					155					160					
gaa	gaa	caa	atc	cta	agg	gca	gct	acg	tcg	atc	tac	ggg	gct	cca	gga	8625
Glu	Glu	Gln	Ile	Leu	Arg	Ala	Ala	Thr	Ser	Ile	Tyr	Gly	Ala	Pro	Gly	
165					170					175					180	
cag	gca	gag	cca	ccc	caa	gct	ttc	ata	gac	gaa	gtt	gcc	aaa	gtc	tat	8673
Gln	Ala	Glu	Pro	Pro	Gln	Ala	Phe	Ile	Asp	Glu	Val	Ala	Lys	Val	Tyr	
			185													

cca aag ccc aag cca aaa ccc aat gct cca aca cag aga ccc cct ggt Pro Lys Pro Lys Pro Lys Pro Asn Ala Pro Thr Gln Arg Pro Pro Gly 230 235 240	8817
cgg ctg ggc cgc tgg atc agg acc gtc tct gat gag gac ctt gag gga Arg Leu Gly Arg Trp Ile Arg Thr Val Ser Asp Glu Asp Leu Glu Gly 245 250 255 260	8865
tcc atc gcc acc atg gtg agc aag ggc gag gag ctg ttc acc ggg gtg Ser Ile Ala Thr Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val 265 270 275	8913
gtg ccc atc ctg gtc gag ctg gac ggc gac gta aac ggc cac aag ttc Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe 280 285 290	8961
agc gtg tcc ggc gag ggc gag ggc gat gcc acc tac ggc aag ctg acc Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr 295 300 305	9009
ctg aag ttc atc tgc acc acc ggc aag ctg ccc gtg ccc tgg ccc acc Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr 310 315 320	9057
ctc gtg acc acc ctg acc tac ggc gtg cag tgc ttc agc cgc tac ccc Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro 325 330 335 340	9105
gac cac atg aag cag cac gac ttc ttc aag tcc gcc atg ccc gaa ggc Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly 345 350 355	9153
tac gtc cag gag cgc acc atc ttc ttc aag gac gac ggc aac tac aag Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys 360 365 370	9201
acc cgc gcc gag gtg aag ttc gag ggc gac acc ctg gtg aac cgc atc Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile 375 380 385	9249
gag ctg aag ggc atc gac ttc aag gag gac ggc aac atc ctg ggg cac Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His 390 395 400	9297
aag ctg gag tac aac tac aac agc cac aac gtc tat atc atg gcc gac Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp 405 410 415 420	9345
aag cag aag aac ggc atc aag gtg aac ttc aag atc cgc cac aac atc Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile 425 430 435	9393
gag gac ggc agc gtg cag ctc gcc gac cac tac cag cag aac acc ccc Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro 440 445 450	9441
atc ggc gac ggc ccc gtg ctg ctg ccc gac aac cac tac ctg agc acc Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr 455 460 465	9489

cag tcc gcc ctg agc aaa gac ccc aac gag aag cgc gat cac atg gtc 9537
 Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val
 470 475 480

ctg ctg gag ttc gtg acc gcc gcc ggg atc act ctc ggc atg gac gag 9585
 Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu
 485 490 495 500

ctg tac aag taa agc 9600
 Leu Tyr Lys

<210> 10
 <211> 503
 <212> PRT
 <213> Artificial sequence

<220>
 <223> pVP2-VP3-GFP protein

<400> 10
 Met Ala Ala Ser Glu Phe Lys Glu Thr Pro Glu Leu Glu Ser Ala Val
 1 5 10 15
 Arg Ala Met Glu Ala Ala Ala Asn Val Asp Pro Leu Phe Gln Ser Ala
 20 25 30
 Leu Ser Val Phe Met Trp Leu Glu Glu Asn Gly Ile Val Thr Asp Met
 35 40 45
 Ala Asn Phe Ala Leu Ser Asp Pro Asn Ala His Arg Met Arg Asn Phe
 50 55 60
 Leu Ala Asn Ala Pro Gln Ala Gly Ser Lys Ser Gln Arg Ala Lys Tyr
 65 70 75 80
 Gly Thr Ala Gly Tyr Gly Val Glu Ala Arg Gly Pro Thr Pro Glu Glu
 85 90 95
 Ala Gln Arg Glu Lys Asp Thr Arg Ile Ser Lys Lys Met Glu Thr Met
 100 105 110
 Gly Ile Tyr Phe Ala Thr Pro Glu Trp Val Ala Leu Asn Gly His Arg
 115 120 125
 Gly Pro Ser Pro Gly Gln Val Lys Tyr Trp Gln Asn Lys Arg Glu Ile
 130 135 140
 Pro Asp Pro Asn Glu Asp Tyr Leu Asp Tyr Val His Ala Glu Lys Ser
 145 150 155 160
 Arg Leu Ala Ser Glu Glu Gln Ile Leu Arg Ala Ala Thr Ser Ile Tyr
 165 170 175
 Gly Ala Pro Gly Gln Ala Glu Pro Pro Gln Ala Phe Ile Asp Glu Val
 180 185 190
 Ala Lys Val Tyr Glu Ile Asn His Gly Arg Gly Pro Asn Gln Glu Gln
 195 200 205

Met Lys Asp Leu Leu Leu Thr Ala Met Glu Met Lys His Arg Asn Pro
 210 215 220
 Arg Arg Ala Leu Pro Lys Pro Lys Pro Lys Pro Asn Ala Pro Thr Gln
 225 230 235 240
 Arg Pro Pro Gly Arg Leu Gly Arg Trp Ile Arg Thr Val Ser Asp Glu
 245 250 255
 Asp Leu Glu Gly Ser Ile Ala Thr Met Val Ser Lys Gly Glu Glu Leu
 260 265 270
 Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn
 275 280 285
 Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr
 290 295 300
 Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val
 305 310 315 320
 Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe
 325 330 335
 Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala
 340 345 350
 Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp
 355 360 365
 Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu
 370 375 380
 Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn
 385 390 395 400
 Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr
 405 410 415
 Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile
 420 425 430
 Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln
 435 440 445
 Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His
 450 455 460
 Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg
 465 470 475 480
 Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu
 485 490 495
 Gly Met Asp Glu Leu Tyr Lys
 500

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/000694

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N7/04 A61K39/12 C12N15/62 C07K14/08 C07K19/00
C12N15/87 C12N15/86 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HU Y ET AL: "Chimeric infectious bursal disease virus-like particles expressed in insect cells and purified by immobilized metal affinity chromatography" BIOTECHNOLOGY AND BIOENGINEERING. INCLUDING: SYMPOSIUM BIOTECHNOLOGY IN ENERGY PRODUCTION AND CONSERVATION, JOHN WILEY & SONS. NEW YORK, US, vol. 63, no. 6, 20 June 1999 (1999-06-20), pages 721-729, XP002190336 ISSN: 0006-3592 page 721 - page 724	1-19
X	WO 01/97839 A (RAHAN MERISTEM; STRAM, YEHUDA; ROGEL, ARIE; EDELBAUM, ORIT; SELA, ILAN) 27 December 2001 (2001-12-27) page 5 - page 6; claims 1-20	1-7, 9, 12-19
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

14 June 2005

Date of mailing of the international search report

27/06/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Paresce, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2005/000694

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>FERNÁNDEZ-ARIAS A ET AL: "Expression of ORF A1 of infectious bursal disease virus results in the formation of virus-like particles"</p> <p>JOURNAL OF GENERAL VIROLOGY, SOCIETY FOR GENERAL MICROBIOLOGY, READING, GB, vol. 79, no. part 5, May 1998 (1998-05), pages 1047-1054, XP002218365</p> <p>ISSN: 0022-1317</p> <p>cited in the application</p> <p>page 1049 - page 1053</p>	1-19
Y	<p>MARTINEZ-TORRECUADRADA J L ET AL:</p> <p>"Different Architectures in the Assembly of Infectious Bursal Disease Virus Capsid Proteins Expressed in Insect Cells"</p> <p>VIROLOGY, ACADEMIC PRESS, ORLANDO, US, vol. 278, no. 2,</p> <p>20 December 2000 (2000-12-20), pages 322-331, XP004435746</p> <p>ISSN: 0042-6822</p> <p>cited in the application</p> <p>the whole document</p>	1-19
Y	<p>MARTINEZ-TORRECUADRADA J L ET AL:</p> <p>"Structure-dependent efficacy of infectious bursal disease virus (IBDV) recombinant vaccines"</p> <p>VACCINE, BUTTERWORTH SCIENTIFIC, GUILDFORD, GB,</p> <p>vol. 21, no. 23, 4 July 2003 (2003-07-04), pages 3342-3350, XP004429746</p> <p>ISSN: 0264-410X</p> <p>the whole document</p>	1-19
Y	<p>US 5 788 970 A (VAKHARIA ET AL)</p> <p>4 August 1998 (1998-08-04)</p> <p>cited in the application</p> <p>columns 14-15</p>	1-19
P,X	<p>ONA A ET AL: "The C-terminal domain of the pVP2 precursor is essential for the interaction between VP2 and VP3, the capsid polypeptides of infectious bursal disease virus"</p> <p>VIROLOGY, ACADEMIC PRESS, ORLANDO, US, vol. 322, no. 1,</p> <p>25 April 2004 (2004-04-25), pages 135-142, XP004500338</p> <p>ISSN: 0042-6822</p> <p>page 136 - page 137</p>	1-10, 12-19

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP2005/000694

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0197839	A	27-12-2001	WO 0197839 A1	27-12-2001
			AU 5244900 A	02-01-2002
			EP 1296710 A1	02-04-2003
<hr/>				
US 5788970	A	04-08-1998	AU 696656 B2	17-09-1998
			AU 2129195 A	17-10-1995
			CA 2186856 A1	05-10-1995
			EP 0755259 A1	29-01-1997
			JP 9510873 T	04-11-1997
			US 6017759 A	25-01-2000
			WO 9526196 A1	05-10-1995
			US 6156314 A	05-12-2000
<hr/>				

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.